

New England Biolabs Certificate of Analysis

Product Name: *Nuclease-free Water*
Catalog Number: *B1500L*
Packaging Lot Number: *10160920*
Expiration Date: *02/2024*
Storage Temperature: *25°C*
Specification Version: *PS-B1500S/L v2.0*

| Nuclease-free Water Component List | | | |
|------------------------------------|-----------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| B1500SVIAL | Nuclease-free Water | 10131673 | Pass |

| Assay Name/Specification | Lot # 10160920 |
|--|----------------|
| <p>RNase Activity (Extended Digestion, Water) A 10 µl reaction in 1X NEBuffer 4 containing 40 ng of RNA transcript with Nuclease-free Water is incubated at 37°C. After incubation for 16 hours, no detectable degradation of the RNA is observed as determined by gel electrophoresis using fluorescent detection.</p> | Pass |
| <p>UV-Visible Scan A UV-Visible scan using a spectrophotometer that covers the range of 200nm to 800nm will have no detectable peaks above background.</p> | Pass |
| <p>qPCR DNA Contamination (E. coli Genomic, Water) Nuclease-free Water is used to make a qPCR master mix and screened across a 96 well plate for the presence of E. coli genomic DNA using 40 cycles of SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Melt curve analysis results in < 5% positive samples above background.</p> | Pass |
| <p>Endonuclease Activity (Nicking, Water) A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 RF I DNA with Nuclease-free Water incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p> | Pass |
| <p>Endotoxin Testing (Endosafe®) Each test channel of the cartridge is loaded with 25 µl of Nuclease-free Water, then placed into the Endosafe MCS reader for analysis resulting in a measurement of <0.01 EU/ml.</p> | Pass |


| Assay Name/Specification | Lot # 10160920 |
|---|--------------------|
| <p>Non-Specific DNase Activity (16 Hour, Water) A 50 µl reaction in CutSmart[®] Buffer containing 1 µg of PhiX174-HaeIII DNA with Nuclease-free Water incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p> | <p>Pass</p> |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Christie Vazquez
Production Scientist
05 Aug 2022



Nick Privitera
Packaging Quality Control Inspector
05 Aug 2022