

Alkaline Phosphatase, Calf Intestinal (CIP)



1-800-632-7799
info@neb.com
www.neb.com



M0290S 065141116111

M0290S



1,000 units Lot: 0651411 Exp: 11/16

10,000 U/ml Store at -20°C

Description: Alkaline Phosphatase, Calf Intestinal (CIP) nonspecifically catalyzes the dephosphorylation of 5' and 3' ends of DNA and RNA phosphomonoesters. Also, CIP hydrolyses ribo-, as well as deoxyribonucleoside triphosphates (NTPs and dNTPs). CIP is useful in many molecular biology applications such as the removal of phosphorylated ends of DNA and RNA for subsequent use in cloning or end-labeling of probes. In cloning, dephosphorylation prevents

Now with CutSmart Reaction Buffer

Alkaline Phosphatase, Calf Intestinal (CIP)



1-800-632-7799
info@neb.com
www.neb.com



M0290S 065141116111

M0290S



1,000 units Lot: 0651411 Exp: 11/16

10,000 U/ml Store at -20°C

Description: Alkaline Phosphatase, Calf Intestinal (CIP) nonspecifically catalyzes the dephosphorylation of 5' and 3' ends of DNA and RNA phosphomonoesters. Also, CIP hydrolyses ribo-, as well as deoxyribonucleoside triphosphates (NTPs and dNTPs). CIP is useful in many molecular biology applications such as the removal of phosphorylated ends of DNA and RNA for subsequent use in cloning or end-labeling of probes. In cloning, dephosphorylation prevents

Now with CutSmart Reaction Buffer

religation of linearized plasmid DNA. The enzyme acts on 5' protruding, 5' recessed and blunt ends. CIP may also be used to degrade unincorporated dNTPs in PCR reactions to prepare templates for DNA sequencing or SNP analysis.

Source: Calf intestinal mucosa

Specific Activity: ~ 3,000 units/mg

Molecular Weight: CIP is a homodimer. The molecular weight of the monomer is 69 kDa.

Applications:

- Dephosphorylation of cloning vector DNA to prevent recircularization during ligation
- Dephosphorylation of DNA prior to end-labeling using T4 Polynucleotide Kinase
- Treatment of dNTPs in PCR reactions prior to sequencing or SNP analysis
- Dephosphorylation of DNA and RNA

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 8.2 @ 25°C), 1 mM MgCl₂, 0.1 mM ZnCl₂ and 50% glycerol.

religation of linearized plasmid DNA. The enzyme acts on 5' protruding, 5' recessed and blunt ends. CIP may also be used to degrade unincorporated dNTPs in PCR reactions to prepare templates for DNA sequencing or SNP analysis.

Source: Calf intestinal mucosa

Specific Activity: ~ 3,000 units/mg

Molecular Weight: CIP is a homodimer. The molecular weight of the monomer is 69 kDa.

Applications:

- Dephosphorylation of cloning vector DNA to prevent recircularization during ligation
- Dephosphorylation of DNA prior to end-labeling using T4 Polynucleotide Kinase
- Treatment of dNTPs in PCR reactions prior to sequencing or SNP analysis
- Dephosphorylation of DNA and RNA

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 8.2 @ 25°C), 1 mM MgCl₂, 0.1 mM ZnCl₂ and 50% glycerol.

Reagents Supplied with Enzyme:

10X CutSmart™ Reaction Buffer

Reaction Conditions: 1X CutSmart Reaction Buffer.

Incubate at 37°C.

1X CutSmart Reaction Buffer:

50 mM Potassium acetate

20 mM Tris-acetate

10 mM Magnesium acetate

100 µg/ml BSA

pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme that hydrolyzes 1 µmol of *p*-Nitrophenyl Phosphate, PNPP (NEB #P0757) in a total reaction volume of 1 ml in 1 minute at 37°C.

Unit Assay Conditions: 1 M diethanolamine-HCl (pH 9.8), 0.5 mM MgCl₂, 50 mM *p*-nitrophenylphosphate and enzyme. These conditions are only used for quantitating enzyme activity.

Reagents Supplied with Enzyme:

10X CutSmart™ Reaction Buffer

Reaction Conditions: 1X CutSmart Reaction Buffer.

Incubate at 37°C.

1X CutSmart Reaction Buffer:

50 mM Potassium acetate

20 mM Tris-acetate

10 mM Magnesium acetate

100 µg/ml BSA

pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme that hydrolyzes 1 µmol of *p*-Nitrophenyl Phosphate, PNPP (NEB #P0757) in a total reaction volume of 1 ml in 1 minute at 37°C.

Unit Assay Conditions: 1 M diethanolamine-HCl (pH 9.8), 0.5 mM MgCl₂, 50 mM *p*-nitrophenylphosphate and enzyme. These conditions are only used for quantitating enzyme activity.

Functional Assay: Dephosphorylation with CIP of a restriction enzyme-digested vector DNA with 5' recessed ends, the least favorable type for dephosphorylation, reduces re-ligation to < 0.5% compared to untreated control as measured by transformation into *E. coli*. CIP has been functionally tested in the following protocol:

Protocol for Dephosphorylation of 5'-ends of DNA using CIP

1. Prepare a 20 µl reaction as follows:

| | |
|----------------------------|---------------------|
| DNA | 1 pmol of DNA ends* |
| CutSmart Buffer (10X) | 2 µl |
| CIP | 1 unit |
| H ₂ O, purified | to 20 µl** |

2. Incubate at 37°C for 30 minutes.

*Note: 1 pmol of DNA ends is about 1 µg of a 3 kb plasmid.

**Scale larger reaction volumes proportionally.

3. Purify DNA by gel purification, spin-column or phenol extraction.

CERTIFICATE OF ANALYSIS

Functional Assay: Dephosphorylation with CIP of a restriction enzyme-digested vector DNA with 5' recessed ends, the least favorable type for dephosphorylation, reduces re-ligation to < 0.5% compared to untreated control as measured by transformation into *E. coli*. CIP has been functionally tested in the following protocol:

Protocol for Dephosphorylation of 5'-ends of DNA using CIP

1. Prepare a 20 µl reaction as follows:

| | |
|----------------------------|---------------------|
| DNA | 1 pmol of DNA ends* |
| CutSmart Buffer (10X) | 2 µl |
| CIP | 1 unit |
| H ₂ O, purified | to 20 µl** |

2. Incubate at 37°C for 30 minutes.

*Note: 1 pmol of DNA ends is about 1 µg of a 3 kb plasmid.

**Scale larger reaction volumes proportionally.

3. Purify DNA by gel purification, spin-column or phenol extraction.

CERTIFICATE OF ANALYSIS

Protocol for Dephosphorylation of 5'-ends of DNA using CIP in Restriction Enzyme Reaction

1. Digest 1–5 µg of plasmid DNA in a 20 µl reaction as follows:

| | |
|---------------------------------|----------|
| DNA | ≥ 1 µl |
| Restriction Enzyme Buffer (10X) | 2 µl |
| Restriction Endonuclease | 1 µl |
| H ₂ O, purified | to 20 µl |

Note: Scale larger reaction volumes proportionally.

2. Incubate at 37°C for 60 minutes or follow manufacturer's recommendations.
3. Add 1 unit of CIP for every 1 pmol of DNA ends (about 1 µg of a 3 kb plasmid) and incubate at 37°C for 30–60 minutes.
4. Purify DNA by gel purification, spin-column or phenol extraction.
5. Proceed with ligation.

Usage Notes:

1. CIP, as are most alkaline phosphatases, is a Zn²⁺ and Mg²⁺-dependent enzyme. Our formulation of its storage buffer provides Zn²⁺ and Mg²⁺, which does not require supplemental zinc or other additives in reactions with CIP.

Protocol for Dephosphorylation of 5'-ends of DNA using CIP in Restriction Enzyme Reaction

1. Digest 1–5 µg of plasmid DNA in a 20 µl reaction as follows:

| | |
|---------------------------------|----------|
| DNA | ≥ 1 µl |
| Restriction Enzyme Buffer (10X) | 2 µl |
| Restriction Endonuclease | 1 µl |
| H ₂ O, purified | to 20 µl |

Note: Scale larger reaction volumes proportionally.

2. Incubate at 37°C for 60 minutes or follow manufacturer's recommendations.
3. Add 1 unit of CIP for every 1 pmol of DNA ends (about 1 µg of a 3 kb plasmid) and incubate at 37°C for 30–60 minutes.
4. Purify DNA by gel purification, spin-column or phenol extraction.
5. Proceed with ligation.

Usage Notes:

1. CIP, as are most alkaline phosphatases, is a Zn²⁺ and Mg²⁺-dependent enzyme. Our formulation of its storage buffer provides Zn²⁺ and Mg²⁺, which does not require supplemental zinc or other additives in reactions with CIP.

2. CIP is also active in 1X NEBuffers 1.1, 2.1, 3.1, as well as NEBuffers 1, 2, 3, 4 and unique NEBuffer for EcoRI.
3. CIP activity is enhanced in the presence of monovalent salts.
4. CIP is inhibited by metal chelators (e.g. EDTA), inorganic phosphate and phosphate analogs.

Quality Controls Assays

Exonuclease Activity: Incubation of a 50 µl reaction containing 50 units of CIP with 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA for 4 hours at 37°C released < 0.1% of the total radioactivity.

Endonuclease Activity: Incubation of a 50 µl reaction containing 50 units of CIP with 1 µg of φX174 RF I DNA for 4 hours at 37°C resulted in < 10% conversion to RF II as determined by agarose gel electrophoresis.

RNase Activity: Incubation of a 10 µl reaction containing 10 units of CIP with 40 ng of fluorescein labeled RNA transcript for 4 hours at 37°C resulted in < 10% degradation of the RNA as determined by gel electrophoresis using fluorescence detection.

2. CIP is also active in 1X NEBuffers 1.1, 2.1, 3.1, as well as NEBuffers 1, 2, 3, 4 and unique NEBuffer for EcoRI.
3. CIP activity is enhanced in the presence of monovalent salts.
4. CIP is inhibited by metal chelators (e.g. EDTA), inorganic phosphate and phosphate analogs.

Quality Controls Assays

Exonuclease Activity: Incubation of a 50 µl reaction containing 50 units of CIP with 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA for 4 hours at 37°C released < 0.1% of the total radioactivity.

Endonuclease Activity: Incubation of a 50 µl reaction containing 50 units of CIP with 1 µg of φX174 RF I DNA for 4 hours at 37°C resulted in < 10% conversion to RF II as determined by agarose gel electrophoresis.

RNase Activity: Incubation of a 10 µl reaction containing 10 units of CIP with 40 ng of fluorescein labeled RNA transcript for 4 hours at 37°C resulted in < 10% degradation of the RNA as determined by gel electrophoresis using fluorescence detection.

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

Heat Inactivation: No

References:

1. Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, (2nd ed.), (p. 5.72). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
2. Mossner, E., Boll, M. and Pfeleiderer, G. (1980) *Hoppe-Seyler's Z. Physiol. Chem.* 361, 543–549.

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

Heat Inactivation: No

References:

1. Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, (2nd ed.), (p. 5.72). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
2. Mossner, E., Boll, M. and Pfeleiderer, G. (1980) *Hoppe-Seyler's Z. Physiol. Chem.* 361, 543–549.

Companion Products Sold Separately:

| | |
|---------------|---------------|
| T4 DNA Ligase | |
| #M0202S | 20,000 units |
| #M0202L | 100,000 units |
| #M0202T | 20,000 units |
| #M0202M | 100,000 units |

| | |
|---------------------|----------|
| Quick Ligation™ Kit | |
| #M2200S | 30 rxns |
| #M2200L | 150 rxns |

| | |
|--------------------------------------|----------|
| Instant Sticky-end Ligase Master Mix | |
| #M0370S | 50 rxns |
| #M0370L | 250 rxns |

| | |
|----------------------------|----------|
| Blunt/TA Ligase Master Mix | |
| #M0367S | 50 rxns |
| #M0367L | 250 rxns |



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

CUTSMART™ and QUICK LIGATION™ are trademarks of New England Biolabs, Inc.

Companion Products Sold Separately:

| | |
|---------------|---------------|
| T4 DNA Ligase | |
| #M0202S | 20,000 units |
| #M0202L | 100,000 units |
| #M0202T | 20,000 units |
| #M0202M | 100,000 units |

| | |
|---------------------|----------|
| Quick Ligation™ Kit | |
| #M2200S | 30 rxns |
| #M2200L | 150 rxns |

| | |
|--------------------------------------|----------|
| Instant Sticky-end Ligase Master Mix | |
| #M0370S | 50 rxns |
| #M0370L | 250 rxns |

| | |
|----------------------------|----------|
| Blunt/TA Ligase Master Mix | |
| #M0367S | 50 rxns |
| #M0367L | 250 rxns |



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

CUTSMART™ and QUICK LIGATION™ are trademarks of New England Biolabs, Inc.