

## New England Biolabs Certificate of Analysis


**Product Name:** Isothermal Amplification Buffer  
**Catalog Number:** B0537S  
**Concentration:** 10 X Concentrate  
**Lot Number:** 10038442  
**Expiration Date:** 02/2022  
**Storage Temperature:** -20°C  
**Specification Version:** PS-B0537S v1.0  
**Composition (1X):** 20 mM Tris-HCl, 50 mM KCl, 10 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 2 mM MgSO<sub>4</sub>, 0.1 % Tween® 20, (pH 8.8 @ 25°C)

Isothermal Amplification Buffer Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
B0537SVIAL	Isothermal Amplification Buffer	10035085	Pass

Assay Name/Specification	Lot # 10038442
<b>Endonuclease Activity (Nicking, Buffer)</b> A 50 µl reaction in 2X Isothermal Amplification Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Non-Specific DNase Activity (16 hour, Buffer)</b> A 50 µl reaction in 2X Isothermal Amplification Buffer containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>pH (buffers/solutions)</b> The pH of 10X Isothermal Amplification Buffer is between pH 8.7 and 8.9 at 25°C.	Pass
<b>Phosphatase Activity (pNPP, Buffer)</b> A 200 µl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl <sub>2</sub> containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 40 µl Isothermal Amplification Buffer incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
<b>qPCR DNA Contamination (E. coli Genomic, Buffer)</b> A minimum of 1 µl of Isothermal Amplification Buffer is screened for the presence of	Pass

Assay Name/Specification	Lot # 10038442
<p>E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is <math>\leq 1</math> E. coli genome.</p> <p><b>RNase Activity Assay (4 Hour Digestion)</b> A 10 <math>\mu</math>l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 <math>\mu</math>l of Isothermal Amplification Buffer is incubated at 37°C. After incubation for 4 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.



Doreen Duquette  
Production Scientist  
21 Feb 2019



Michael Tonello  
Packaging Quality Control Inspector  
20 Mar 2019