

## New England Biolabs Certificate of Analysis

**Product Name:** NEBuffer™ 3.1  
**Catalog Number:** B7203S  
**Concentration:** 10 X Concentrate  
**Packaging Lot Number:** 10082863  
**Expiration Date:** 06/2023  
**Storage Temperature:** -20°C  
**Specification Version:** PS-B7203S v1.0  
**Composition (1X):** 100 mM NaCl, 50 mM Tris-HCl, 10 mM MgCl<sub>2</sub>, 100 µg/ml BSA, (pH 7.9 @ 25°C)


NEBuffer™ 3.1 Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
B7203SVIAL	NEBuffer™ 3.1	10077593	Pass

Assay Name/Specification	Lot # 10082863
<b>Conductivity (buffers/solutions)</b> The conductivity of 10X NEBuffer 3.1 is between 84 and 101 mS at 25°C.	Pass
<b>Endonuclease Activity (Nicking, Buffer)</b> A 50 µl reaction in 1X NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Functional Testing (Restriction Digest, Buffer)</b> A 50 µl reaction in 1X NEBuffer 3.1 containing 1 µg of pBC4 DNA and 1 unit of NotI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
<b>Functional Testing (Restriction Digest, Buffer)</b> A 50 µl reaction in 1X NEBuffer 3.1 containing 1 µg of Lambda DNA and 1 unit of AseI incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	Pass
<b>Non-Specific DNase Activity (16 hour, Buffer)</b> A 50 µl reaction in 1X NEBuffer 3.1 containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass

Assay Name/Specification	Lot # 10082863
<p><b>pH (buffers/solutions)</b> The pH of 10X NEBuffer 3.1 is between pH 7.8 and 8.0 at 25°C.</p>	<b>Pass</b>
<p><b>RNase Activity (Buffer)</b> A 10 µl reaction in 1X NEBuffer 3.1 containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, &gt;90% of the substrate RNA remains intact as determined by fluorescent detection.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.




---

Doreen Duquette  
Production Scientist  
13 Jul 2020




---

Josh Hersey  
Packaging Quality Control Inspector  
19 Aug 2020