

New England Biolabs Certificate of Analysis

Product Name: Standard Taq (Mg-free) Reaction Buffer Pack
 Catalog Number: B9015S
 Concentration: 10 X Concentrate
 Packaging Lot Number: 10055983
 Expiration Date: 01/2024
 Storage Temperature: -20°C
 Specification Version: PS-B9015S v1.0
 Composition (1X): 10 mM Tris-HCl, 50 mM KCl, (pH 8.3 @ 25°C)

Standard Taq (Mg-free) Reaction Buffer Pack Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
B9021SVIAL	Magnesium Chloride (MgCl ₂) Solution	10054951	Pass
B9015SVIAL	Standard Taq (Mg-free) Reaction Buffer Pack	10033771	Pass


Assay Name/Specification	Lot # 10055983
<p>qPCR DNA Contamination (E. coli Genomic, Buffer) A minimum of 1 µl of Standard Taq (Mg-free) Reaction Buffer is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass
<p>Non-Specific DNase Activity (16 hour, Mg-Free Buffer) A 50 µl reaction in 2X Standard Taq (Mg-free) Reaction Buffer and 3 mM MgCl₂ containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>PCR Amplification (5 kb Lambda DNA, Mg-Free Buffer) A 50 µl reaction in Standard Taq (Mg-free) Reaction Buffer and 1.5 mM MgCl₂ in the presence of 200 µM dNTPs and 0.2 µM primers containing 5 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 5 kb product.</p>	Pass
<p>Phosphatase Activity (pNPP, Buffer) A 200 µl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 40 µl Standard Taq (Mg-free)</p>	Pass

Assay Name/Specification	Lot # 10055983
<p>Reaction Buffer incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	
<p>pH (buffers/solutions) The pH of 10X Standard Taq (Mg-free) Reaction Buffer is between pH 8.2 and 8.4 at 25°C.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Standard Taq (Mg-free) Reaction Buffer is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>Endonuclease Activity (Nicking, Mg-Free Buffer) A 50 µl reaction in 2X Standard Taq (Mg-free) Reaction Buffer and 3 mM MgCl₂ containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Christie Vazquez
Production Scientist
05 Feb 2019



Michael Tonello
Packaging Quality Control Inspector
31 Dec 2019