

New England Biolabs Certificate of Analysis

Product Name: OneTaq[®] GC Reaction Buffer Pack
Catalog #: B9023S
Concentration: 5X Concentrate
Lot #: 0031708
Assay Date: 08/2017
Expiration Date: 8/2020
Storage Temp: -20°C
Composition (1X): 80 mM Tris-SO₄, 20 mM (NH₄)₂SO₄, 2 mM MgSO₄, 5 % Glycerol, 5 % DMSO, 0.06 % IGEPAL[®] CA-630, 0.05 % Tween[®] 20, (pH 9.2 @ 25°C)
Specification Version: PS-B9023S v1.0
Effective Date: 02 Aug 2017

Assay Name/Specification (minimum release criteria)	Lot #0031708
Endonuclease Activity (Nicking, Buffer) - A 50 µl reaction in 2X OneTaq [®] GC Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 hour, Buffer) - A 50 µl reaction in 2X OneTaq [®] GC Reaction Buffer containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
PCR Amplification (Buffer Dependent, >65% GC-rich, Buffer) - A 25 µl reaction in OneTaq [®] GC Reaction Buffer in the presence of 200 µM dNTPs and 0.2 µM primers containing 10 ng Human Genomic DNA with 0.625 units of OneTaq [®] DNA Polymerase for 30 cycles of PCR amplification results in the buffer-dependent production of the expected 737 bp product.	Pass
PCR Amplification (Enhancer Dependent, >70% GC-rich, Buffer) - A 25 µl reaction in OneTaq [®] GC Reaction Buffer and 20% OneTaq [®] High GC Enhancer in the presence of 200 µM dNTPs and 0.2 µM primers containing 10 ng Human Genomic DNA with 0.625 units of OneTaq [®] DNA Polymerase for 30 cycles of PCR amplification results in the enhancer-dependent production of the expected 627 bp product.	Pass
pH (buffers/solutions) - The pH of 5X OneTaq [®] GC Reaction Buffer is between pH 9.1 and 9.3 at 25°C.	Pass



New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0031708
<p>Phosphatase Activity (pNPP, Buffer) - A 200 µl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl₂ containing 2.5 mM <i>p</i>-Nitrophenyl Phosphate (pNPP) and a minimum of 80 µl OneTaq[®] GC Reaction Buffer incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	Pass
<p>qPCR DNA Contamination (<i>E. coli</i> Genomic, Buffer) - A minimum of 1 µl of OneTaq[®] GC Reaction Buffer is screened for the presence of <i>E. coli</i> genomic DNA using SYBR[®] Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is ≤ 1 <i>E. coli</i> genome.</p>	Pass
<p>RNase Activity (Extended Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of OneTaq[®] GC Reaction Buffer is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass



Authorized by
Lynne Apone
02 Aug 2017



Inspected by
Tony Spear-Alfonso
05 Sep 2017

