

New England Biolabs Certificate of Analysis

Product Name: SHuffle[®] T7 Express lysY Competent E.coli
Catalog Number: C3030J
Lot Number: 10041121
Expiration Date: 03/2020
Storage Temperature: -80°C
Specification Version: PS-C3030J v1.0

SHuffle [®] T7 Express lysY Competent E.coli Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
C3030JVIAL	SHuffle [®] T7 Express lysY Competent E.coli	10015241	Pass


Assay Name/Specification	Lot # 10041121
Antibiotic Resistance (Chloramphenicol) 15 µl of untransformed Shuffle [®] T7 Express lysY Competent E. coli streaked onto a LB or Rich Broth plate containing Chloramphenicol will form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Resistance (Nitrofurantoin) 15 µl of untransformed Shuffle [®] T7 Express lysY Competent E. coli streaked onto a LB or Rich Broth plate containing Nitrofurantoin will form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Resistance (Spectinomycin) 15 µl of untransformed Shuffle [®] T7 Express lysY Competent E. coli streaked onto a LB or Rich Broth plate containing Spectinomycin will form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Resistance (Streptomycin) 15 µl of untransformed Shuffle [®] T7 Express lysY Competent E. coli streaked onto a LB or Rich Broth plate containing Streptomycin will form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Ampicillin) 15 µl of untransformed Shuffle [®] T7 Express lysY Competent E. coli streaked onto a LB or Rich Broth plate containing Ampicillin will not form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Kanamycin)	Pass

Assay Name/Specification	Lot # 10041121
<p>15 µl of untransformed Shuffle® T7 Express lysY Competent E. coli streaked onto a LB or Rich Broth plate containing Kanamycin will not form colonies after incubation for 16 hours at 37°C.</p>	
<p>Antibiotic Sensitivity (Tetracycline) 15 µl of untransformed Shuffle® T7 Express lysY Competent E. coli streaked onto a LB or Rich Broth plate containing Tetracycline will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Functional Testing (Disulfide Bond Formation) The nuclease NucA requires disulfide bonds for its stability. When expressed at 37°C in E. coli, NucA is toxic to cells only in its oxidized disulfide-bonded state. Transformation of SHuffle® T7 Express lysY Competent E. coli using 100 pg of plasmid that expresses a MBP-NucA fusion results in < 1% of the colonies when compared to a control transformation of its wild type parent strain NEB 10-beta.</p>	Pass
<p>Phage Resistance (φ 80) 15 µl of untransformed Shuffle® T7 Express lysY Competent E. coli streaked onto a Rich Broth plate does not support plaque formation by phage φ 80 after incubation for 16 hours at 37°C.</p>	Pass
<p>Transformation Efficiency 50 µl of SHuffle® T7 Express lysY Competent E. coli cells were transformed with 100 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB-Ampicillin plates at 37°C resulted in >1 x 10⁷ cfu/µg of DNA.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Derek Robinson
Production Scientist
16 Jul 2018



Corey Rabeau
Packaging Quality Control Inspector
21 Mar 2019