

New England Biolabs Certificate of Analysis

Product Name: *Thermostable Inorganic Pyrophosphatase*
Catalog Number: *M0296S*
Concentration: *2,000 U/ml*
Unit Definition: *One unit is the amount of enzyme that will generate 1 μmol of phosphate per minute from inorganic pyrophosphate under standard reaction conditions (a 10 minute reaction at 75°C in 50 mM Tricine [pH 8.5], 1 mM MgCl₂, 0.32 mM PPI, reaction volume of 0.5 ml).*
Packaging Lot Number: *10095241*
Expiration Date: *01/2023*
Storage Temperature: *-20°C*
Storage Conditions: *20 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, (pH 8.0 @ 25°C)*
Specification Version: *PS-M0296S/L v1.0*


Thermostable Inorganic Pyrophosphatase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0296SVIAL	Thermostable Inorganic Pyrophosphatase	10095240	Pass

Assay Name/Specification	Lot # 10095241
<p>Phosphatase Activity (pNPP) A 1 ml reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 10 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units of Thermostable Inorganic Pyrophosphatase incubated for 30 minutes at 75°C yields <0.00001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 μl reaction in NEBuffer 1 containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of Thermostable Inorganic Pyrophosphatase incubated for 4 hours at 75°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 2 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 100 units of Thermostable Inorganic Pyrophosphatase incubated for 4 hours at 75°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass

Assay Name/Specification	Lot # 10095241
dNTPase Activity A 500 µl reaction in CircumVent™ Sequencing Buffer in the presence of 200 µM each dNTPs and a minimum of 100 units Thermostable Inorganic Pyrophosphatase incubated for 1 hour at 75°C results in <0.01 µmole of inorganic phosphate from dNTPs as determined by the AAM assay.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Timothy Meixsell
Production Scientist
27 Jan 2021



Michael Tonello
Packaging Quality Control Inspector
27 Jan 2021