

## New England Biolabs Certificate of Analysis

**Product Name:** T7 DNA Ligase  
**Catalog Number:** M0318L  
**Concentration:** 3,000,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to give 50% ligation of 100 ng of Lambda-HindIII fragments in 30 minutes at 25°C.  
**Packaging Lot Number:** 10088330  
**Expiration Date:** 06/2022  
**Storage Temperature:** -20°C  
**Storage Conditions:** 10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)  
**Specification Version:** PS-M0318S/L v2.0


T7 DNA Ligase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0318LVIAL	T7 DNA Ligase	10077406	Pass
B0535AVIAL	StickTogether™ DNA Ligase Buffer	10077408	Pass

Assay Name/Specification	Lot # 10088330
<p><b>qPCR DNA Contamination (E. coli Genomic)</b>            A minimum of 3000 units of T7 DNA Ligase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass
<p><b>Ligation and Recutting (Terminal Integrity, Digested DNA)</b>            A 20 µl reaction in 1X StickTogether™ DNA Ligase Buffer containing 2 µg of Lambda DNA-HindIII Digest and a minimum of 3,000 units of T7 DNA Ligase incubated for 16 hours at 37°C results in &gt;95% ligation of the DNA fragments as determined by agarose gel electrophoresis. Of these ligated fragments, &gt;95% can be recut with HindIII.</p>	Pass
<p><b>Endonuclease Activity (Nicking)</b>            A 50 µl reaction in NEBuffer 1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 15000 units of T7 DNA Ligase incubated for 4 hours at 37°C results in &lt;10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass

Assay Name/Specification	Lot # 10088330
<p><b>Protein Concentration (A280)</b> The concentration of T7 DNA Ligase is 1 mg/ml +/- 10% as determined by UV absorption at 280 nm. Protein concentration is determined by the Pace method using the extinction coefficient of 69,620 and molecular weight of 41,133 daltons for T7 DNA Ligase (Pace, C.N. et al. (1995) Protein Sci., 4, 2411-2423).</p>	<b>Pass</b>
<p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer 1 containing 1 µg of CIP-treated Lambda-HindIII DNA and a minimum of 3000 units of T7 DNA Ligase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in NEBuffer 1 containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 15000 units of T7 DNA Ligase incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p>	<b>Pass</b>
<p><b>RNase Activity (Extended Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of T7 DNA Ligase is incubated at 37°C. After incubation for 16 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	<b>Pass</b>
<p><b>Protein Purity Assay (SDS-PAGE)</b> T7 DNA Ligase is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Ana Egana  
Production Scientist  
11 Dec 2020



Josh Hersey  
Packaging Quality Control Inspector  
11 Dec 2020