

## New England Biolabs Certificate of Analysis

**Product Name:** *RtcB Ligase*  
**Catalog Number:** *M0458S*  
**Concentration:** *15 µM*  
**Packaging Lot Number:** *10123990*  
**Expiration Date:** *08/2023*  
**Storage Temperature:** *-20°C*  
**Storage Conditions:** *50 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, (pH 7.4 @ 25°C)*  
**Specification Version:** *PS-M0458S v1.0*

RtcB Ligase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
N2080AVIAL	GTP	10117592	Pass
M0458SVIAL	RtcB Ligase	10115449	Pass
B1761SVIAL	10mM MnCl <sub>2</sub>	10117591	Pass
B0790AVIAL	10X RtcB Reaction Buffer	10115451	Pass

Assay Name/Specification	Lot # 10123990
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in RtcB Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 15 pmol of RtcB Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in RtcB Reaction Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 15 pmol of RtcB Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	<b>Pass</b>
<b>Functional Testing (RtcB Ligase)</b> A 20 µl reaction in 1X RtcB Reaction Buffer supplemented with 0.1 mM GTP and 1 mM MnCl <sub>2</sub> containing 0.5 µM of a 17 mer 5' FAM-labeled 3' phosphorylated RNA, 0.5 µM of a 30 mer 5' OH RNA and 0.75 µM RtcB Ligase incubated for 1 hour at 37°C results in ≥ 80% ligation as determined by capillary electrophoresis.	<b>Pass</b>
<b>RNase Activity (Extended Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 15 pmol of RtcB Ligase is incubated at 37°C. After incubation for	<b>Pass</b>

Assay Name/Specification	Lot # 10123990
<p>16 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p> <p><b>Protein Purity Assay (SDS-PAGE)</b> RtcB Ligase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Bhairavi Jani  
Production Scientist  
20 Oct 2021



Michael Tonello  
Packaging Quality Control Inspector  
20 Oct 2021