

## New England Biolabs Certificate of Analysis

**Product Name:** NEBNext<sup>®</sup> FFPE DNA Repair Mix  
**Catalog Number:** M6630L  
**Packaging Lot Number:** 10107940  
**Expiration Date:** 04/2022  
**Storage Temperature:** -20°C  
**Specification Version:** PS-M6630S/L v2.0

| NEBNext <sup>®</sup> FFPE DNA Repair Mix Component List |   |            |                      |
|---|---|------------|----------------------|
| NEB Part Number   | Component Description                       | Lot Number | Individual QC Result |
| M6630LVIAL  | NEBNext <sup>®</sup> FFPE DNA Repair Mix    | 10107941   | Pass                 |
| E6622AAVIAL   | NEBNext <sup>®</sup> FFPE DNA Repair Buffer | 10107942   | Pass                 |

| Assay Name/Specification  | Lot # 10107940 |
|---|----------------|
| <p><b>Functional Testing (Oligonucleotide Cleavage - 8-oxo-guanine)</b><br/>           A 10 µl reaction in ThermoPol<sup>®</sup> Reaction Buffer containing 2.5 pmol of annealed oligo containing 8-oxo-guanine as the non-standard base and 1 µl of the NEBNext<sup>®</sup> FFPE DNA Repair Mix incubated for 1 hour at 37°C resulted in &gt;70% cleavage as determined by polyacrylamide gel electrophoresis.</p>   | <b>Pass</b>    |
| <p><b>Functional Testing (FFPE Repair Mix)</b><br/>           Pretreatment with NEBNext<sup>®</sup> FFPE DNA Repair Mix improves the quality of base calling, especially C &amp; G for FFPE DNA, when compared to an untreated control as determined by sequencing on the Illumina<sup>®</sup> platform. NEBNext<sup>®</sup> FFPE DNA Repair Mix lowers the C:T (same as G:A) mutation for FFPE DNA, which is due to cytosine deamination to U, when compared to an untreated control as determined by sequencing on the Illumina<sup>®</sup> platform.</p> | <b>Pass</b>    |
| <p><b>PCR Amplification (1 kb)</b><br/>           A 48 µl reaction in ThermoPol<sup>®</sup> Reaction Buffer containing 1.5 ng of UV damaged Lambda DNA, 100 µM dNTPs, 500 µM NAD<sup>+</sup> and 1 µl of the NEBNext<sup>®</sup> FFPE DNA Repair Mix was incubated for 15 minutes at 37°C. Addition of 100 µM dNTPs, 0.4 µM L1 primer mix and 2.5 units of Taq DNA Polymerase followed by 25 cycles of PCR resulted in the expected 1 kb specific product.</p>  | <b>Pass</b>    |
| <p><b>Functional Testing (Oligonucleotide Cleavage - Thymine Glycol)</b><br/>           A 10 µl reaction in ThermoPol<sup>®</sup> Reaction Buffer containing 2.5 pmol of annealed oligo containing thymine glycol as the non-standard base and 1 µl of the NEBNext<sup>®</sup> FFPE DNA Repair Mix incubated for 20 minutes at 37°C resulted in &gt;70% cleavage as determined</p>  | <b>Pass</b>    |

| Assay Name/Specification   | Lot # 10107940     |
|--|--------------------|
| <p>by polyacrylamide gel electrophoresis.</p> <p><b>Functional Testing (Oligonucleotide Cleavage - Uracil)</b><br/>A 10 µl reaction in ThermoPol® Reaction Buffer containing 2.5 pmol of annealed oligo containing uracil as the non-standard base and 1 µl of the NEBNext® FFPE DNA Repair Mix incubated for 10 minutes at 37°C resulted in &gt;70% cleavage as determined by polyacrylamide gel electrophoresis.</p> | <p><b>Pass</b></p> |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.




---

Christine Sumner  
Production Scientist  
13 May 2021




---

Michael Tonello  
Packaging Quality Control Inspector  
17 Nov 2021