

## New England Biolabs Certificate of Analysis

**Product Name:** SfaNI  
**Catalog #:** R0172S/L  
**Concentration:** 2,000 units/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of PhiX174 RF I DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Lot #:** 1101602  
**Assay Date:** 02/2016  
**Expiration Date:** 2/2018  
**Storage Temp:** -20°C  
**Storage Buffer:** 300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml BSA  
**Specification Version:** PS-R0172S/L v1.0  
**Effective Date:** 06 Nov 2014

Assay Name/Specification (minimum release criteria)	Lot #1101602
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 6 units of SfaNI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	<b>Pass</b>
<b>Ligation and Recutting (Terminal Integrity)</b> - After a 2-fold over-digestion of PhiX174 DNA with SfaNI, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with SfaNI.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 Hour)</b> - A 50 µl reaction in NEBuffer 3.1 containing 1 µg of PhiX174 DNA and a minimum of 2 Units of SfaNI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	<b>Pass</b>

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



Authorized by  
Derek Robinson  
06 Nov 2014



Inspected by  
Jianying Luo  
27 Jan 2016

