

New England Biolabs Certificate of Analysis


Product Name: *AvrII*
Catalog Number: *R0174S*
Concentration: *5,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (HindIII digest) in 1 hour at 37°C in a total reaction volume of 50 µl.*
Lot Number: *10044758*
Expiration Date: *05/2021*
Storage Temperature: *-20°C*
Storage Conditions: *300 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml BSA*
Specification Version: *PS-R0174S/L v1.0*

AvrII Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0174SVIAL	AvrII	10044760	Pass
B7204SVIAL	CutSmart® Buffer	10042964	Pass
B7024SVIAL	Gel Loading Dye, Purple (6X)	10038711	Pass

Assay Name/Specification	Lot # 10044758
Blue-White Screening (Terminal Integrity) A sample of Litmus28i vector linearized with a 10-fold excess of AvrII, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 Units of AvrII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 50 units of AvrII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 50-fold over-digestion of Lambda HindIII DNA with AvrII, >95% of the DNA	Pass

Assay Name/Specification	Lot # 10044758
<p>fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AvrII.</p> <p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda HindIII DNA and a minimum of 50 Units of AvrII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p> <p>Protein Purity Assay (SDS-PAGE) AvrII is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<p style="text-align: center;">Pass</p> <p style="text-align: center;">Pass</p>

This product has been tested and shown to be in compliance with all specifications.



Doreen Duquette
Production Scientist
12 Apr 2019



Michael Tonello
Packaging Quality Control Inspector
17 May 2019