

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Ncol
Catalog Number: R0193M
Concentration: 50,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10098518
Expiration Date: 02/2023
Storage Temperature: -20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0193T/M v1.0

Ncol Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0193MVIAL	Ncol	10098517	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10089405	Pass	
B6003SVIAL	NEBuffer™ r3.1	10103929	Pass	

Assay Name/Specification	Lot # 10098518
Protein Purity Assay (SDS-PAGE)	Pass
Ncol is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 10 Units of Ncol incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	
g	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and double-stranded [3H] E. coli DNA and a minimum of 50 units of Ncol incubated for 4	
hours at 37°C releases <0.1% of the total radioactivity.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 DNA and a	
minimum of 10 Units of Ncol incubated for 4 hours at 37°C results in <50% conversion to the nicked form as determined by agarose gel electrophoresis.	
to the flicked form as determined by againse gerelectrophoresis.	



R0193M / Lot: 10098518

Page 1 of 2

Assay Name/Specification	Lot # 10098518
Ligation and Recutting (Terminal Integrity) After a 5-fold over-digestion of Lambda DNA with Ncol, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Ncol.	Pass
Blue-White Screening (Terminal Integrity) A sample of LITMUS28i vector linearized with a 10-fold excess of Ncol, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

30 Apr 2021

Josh Hersey

Packaging Quality Control Inspector

30 Apr 2021



R0193M / Lot: 10098518

Page 2 of 2