

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: SgrAl
Catalog Number: R0603L
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of LambdaDNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Lot Number: 10044983
Expiration Date: 05/2021
Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0603S/L v1.0

SgrAl Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0603LVIAL	SgrAI	10044984	Pass	
B7204SVIAL	CutSmart® Buffer	10042783	Pass	

Assay Name/Specification	Lot # 10044983
Non-Specific DNase Activity (16 hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 10 Units of SgrAl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.	Pass
Ligation and Recutting (Terminal Integrity) After a 2-fold over-digestion of Lambda DNA with SgrAI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with SgrAI.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of SgrAl incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking)	Pass



R0603L / Lot: 10044983

Page 1 of 2



Assay Name/Specification	Lot # 10044983
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 30 Units of SgrAl incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

Production Scientist

24 Apr 2019

Michael Tonello

Packaging Quality Control Inspector

15 May 2019

