

New England Biolabs Certificate of Analysis

Product Name: *Esp3I*
Catalog Number: *R0734S*
Concentration: *10,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.*
Packaging Lot Number: *10069704*
Expiration Date: *01/2022*
Storage Temperature: *-20°C*
Storage Conditions: *300 mM NaCl , 10 mM Tris-HCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol , 500 µg/ml BSA, (pH 7.4 @ 25°C)*
Specification Version: *PS-R0734S/L v1.0*

Esp3I Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0734SVIAL	Esp3I	10064300	Pass
B7204SVIAL	CutSmart® Buffer	10071078	Pass
B7024SVIAL	Gel Loading Dye, Purple (6X)	10065746	Pass


Assay Name/Specification	Lot # 10069704
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of Esp3I incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (15 minute Digest) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and 1 µl of Esp3I incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with Esp3I, ≥95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with Esp3I.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of Esp3I incubated for 16 hours at 37°C results in a DNA pattern free of	Pass

Assay Name/Specification	Lot # 10069704
detectable nuclease degradation as determined by agarose gel electrophoresis.	
<p>Protein Purity Assay (SDS-PAGE) Esp3I is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 μl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 10 units of Esp3I is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



JianYing Luo
Production Scientist
11 May 2020



Jay Minichiello
Packaging Quality Control Inspector
11 May 2020