

New England Biolabs Certificate of Analysis

Product Name: *SacI-HF[®]*
Catalog Number: *R3156S*
Concentration: *20,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (HindIII digest) in 1 hour at 37°C in a total reaction volume of 50 µl.*
Lot Number: *10026502*
Expiration Date: *10/2020*
Storage Temperature: *-20°C*
Storage Conditions: *100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA*
Specification Version: *PS-R3156S/L v2.0*

SacI-HF[®] Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R3156SVIAL	SacI-HF [®]	10026500	Pass
B7204SVIAL	CutSmart [®] Buffer	10021122	Pass
B7024SVIAL	Gel Loading Dye, Purple (6X)	10021129	Pass

Assay Name/Specification	Lot # 10026502
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of pXba DNA with SacI-HF [™] , >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with SacI-HF [™] .	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart [™] Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 100 Units of SacI-HF [™] incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) SacI-HF [™] is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
Blue-White Screening (Terminal Integrity) A sample of LITMUS28i vector linearized with a 10-fold excess of SacI-HF [™] ,	Pass

Assay Name/Specification	Lot # 10026502
religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	
<p>Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 60 Units of SacI-HF™ incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of SacI-HF™ incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Anthony Francis
Production Scientist
31 Oct 2018



Michael Tonello
Packaging Quality Control Inspector
29 Nov 2018