

## New England Biolabs Certificate of Analysis

**Product Name:** BbsI-HF®  
**Catalog Number:** R3539S  
**Concentration:** 20,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Lot Number:** 10054974  
**Expiration Date:** 06/2021  
**Storage Temperature:** -20°C  
**Storage Conditions:** 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 500 µg/ml BSA, (pH 7.4 @ 25°C)  
**Specification Version:** PS-R3539S/L v1.0

BbsI-HF® Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R3539SVIAL	BbsI-HF®	10047895	Pass
B7204SVIAL	CutSmart® Buffer	10053981	Pass
B7024SVIAL	Gel Loading Dye, Purple (6X)	10050274	Pass


Assay Name/Specification	Lot # 10054974
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled pUC19 DNA and a minimum of 60 units of BbsI-HF incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of BbsI-HF incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Functional Testing (15 minute Digest)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and 1 µl of BbsI-HF incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of Lambda DNA with BbsI-HF, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	Pass

Assay Name/Specification	Lot # 10054974
<p>&gt;95% can be recut with BbsI-HF.</p> <p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of BbsI-HF incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<p><b>Pass</b></p>
<p><b>Protein Purity Assay (SDS-PAGE)</b> BbsI-HF is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.



Stephanie Cornelio  
Production Scientist  
27 Jun 2019



Jay Minichiello  
Packaging Quality Control Inspector  
25 Sep 2019