

New England Biolabs Certificate of Analysis


Product Name: Agel-HF®
Catalog Number: R3552L
Concentration: 20,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction of 50 µl.
Lot Number: 10036693
Expiration Date: 11/2020
Storage Temperature: -20°C
Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA
Specification Version: PS-R3552S/L v1.0

| Agel-HF® Component List | | | |
|-------------------------|------------------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| R3552LVIAL | Agel-HF® | 10028840 | Pass |
| B7204SVIAL | CutSmart® Buffer | 10031567 | Pass |
| B7024SVIAL | Gel Loading Dye, Purple (6X) | 10021141 | Pass |

| Assay Name/Specification | Lot # 10036693 |
|--|----------------|
| <p>Blue-White Screening (Terminal Integrity) A sample of LITMUS28i vector linearized with a 10-fold excess of Agel-HF™, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.</p> | Pass |
| <p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 200 units of Agel-HF™ incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p> | Pass |
| <p>Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with Agel-HF™, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Agel-HF™.</p> | Pass |
| <p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of Agel-HF™ incubated for 16 hours at 37°C results in a DNA pattern free</p> | Pass |

| Assay Name/Specification | Lot # 10036693 |
|--|----------------|
| of detectable nuclease degradation as determined by agarose gel electrophoresis. | |

This product has been tested and shown to be in compliance with all specifications.



Jianying Luo
Production Scientist
14 Dec 2018



Michael Tonello
Packaging Quality Control Inspector
26 Feb 2019