

T4 RNA Ligase Reaction Buffer



1-800-632-7799
info@neb.com
www.neb.com



B0216L 011150818081

B0216L

3.0 ml **Lot: 0111508** **Exp: 8/18**
10X Concentration **Store at -20°C**

Description: New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X T4 RNA Ligase Reaction Buffer:

50 mM Tris-HCl
10 mM MgCl₂
1 mM DTT
pH 7.5 @ 25°C

Reagents Supplied with Buffer:

10 mM ATP
50% PEG8000

Quality Control Assays

16-Hour Incubation: 1 µg of HaeIII digested φX174 RF I DNA in 50 µl of this reaction buffer at 1X concentration showed no detectable non-specific nuclease degradation after incubation at 37°C for 16 hours.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in a 50 µl reaction resulted in less than 5% conversion to RF II.

RNase Activity: Incubation of 40 ng of RNA transcripts in 50 µl of this reaction buffer at a 1X concentration for 2 hours at 37°C resulted in no detectable degradation of the RNA as determined by gel electrophoresis.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

T4 RNA Ligase Reaction Buffer



1-800-632-7799
info@neb.com
www.neb.com



B0216L 011150818081

B0216L

3.0 ml **Lot: 0111508** **Exp: 8/18**
10X Concentration **Store at -20°C**

Description: New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X T4 RNA Ligase Reaction Buffer:

50 mM Tris-HCl
10 mM MgCl₂
1 mM DTT
pH 7.5 @ 25°C

Reagents Supplied with Buffer:

10 mM ATP
50% PEG8000

Quality Control Assays

16-Hour Incubation: 1 µg of HaeIII digested φX174 RF I DNA in 50 µl of this reaction buffer at 1X concentration showed no detectable non-specific nuclease degradation after incubation at 37°C for 16 hours.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in a 50 µl reaction resulted in less than 5% conversion to RF II.

RNase Activity: Incubation of 40 ng of RNA transcripts in 50 µl of this reaction buffer at a 1X concentration for 2 hours at 37°C resulted in no detectable degradation of the RNA as determined by gel electrophoresis.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS