

## M-MuLV Reverse Transcriptase Reaction Buffer



1-800-632-7799  
info@neb.com  
www.neb.com



B0253S 001160319031

# B0253S

### 6.0 ml M-MuLV Reverse Transcriptase

Reaction Buffer (10X) Lot: 0011603

Store at -20°C Exp: 3/19

**Description:** New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

### 1X M-MuLV Reverse Transcriptase

#### Reaction Buffer:

50 mM Tris-HCl  
75 mM KCl  
3 mM MgCl<sub>2</sub>  
10 mM dithiothreitol  
pH 8.3 @ 25°C

## M-MuLV Reverse Transcriptase Reaction Buffer



1-800-632-7799  
info@neb.com  
www.neb.com



B0253S 001160319031

# B0253S

### 6.0 ml M-MuLV Reverse Transcriptase

Reaction Buffer (10X) Lot: 0011603

Store at -20°C Exp: 3/19

**Description:** New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

### 1X M-MuLV Reverse Transcriptase

#### Reaction Buffer:

50 mM Tris-HCl  
75 mM KCl  
3 mM MgCl<sub>2</sub>  
10 mM dithiothreitol  
pH 8.3 @ 25°C

### Quality Control Assays

**16-Hour Incubation:** 1 µg of HaeIII digested φX174 RF I DNA in 50 µl of this reaction buffer at 1X concentration showed no detectable non-specific nuclease degradation after incubation at 37°C for 16 hours.

**Endonuclease Activity:** Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in a 50 µl reaction resulted in less than 5% conversion to RF II.

**RNase Activity:** Incubation of 40 ng of RNA transcripts in 50 µl of this reaction buffer at a 1X concentration for 2 hours at 37°C resulted in no detectable degradation of the RNA as determined by gel electrophoresis.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

### Quality Control Assays

**16-Hour Incubation:** 1 µg of HaeIII digested φX174 RF I DNA in 50 µl of this reaction buffer at 1X concentration showed no detectable non-specific nuclease degradation after incubation at 37°C for 16 hours.

**Endonuclease Activity:** Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in a 50 µl reaction resulted in less than 5% conversion to RF II.

**RNase Activity:** Incubation of 40 ng of RNA transcripts in 50 µl of this reaction buffer at a 1X concentration for 2 hours at 37°C resulted in no detectable degradation of the RNA as determined by gel electrophoresis.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS