

## Lambda Exonuclease Reaction Buffer



1-800-632-7799  
info@neb.com  
www.neb.com



B0262S 002120915091

# B0262S

**6.0 ml Lambda Exonuclease Reaction Buffer (10X)**

**Store at -20°C Lot: 0021209 Exp: 9/15**

**Description:** New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

### **1X Lambda Exonuclease Reaction Buffer:**

67 mM Glycine-KOH  
2.5 mM MgCl<sub>2</sub>  
50 µg/ml BSA  
pH 9.4 @ 25°C

### **Quality Control Assays**

**16-Hour Incubation:** A 50 µl reaction containing this reaction buffer at a 1X concentration and 1 µg of HaeIII digested φX174 RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

**Endonuclease Activity:** Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reactions resulted in less than 5% conversion to RF II.

CERTIFICATE OF ANALYSIS

## Lambda Exonuclease Reaction Buffer



1-800-632-7799  
info@neb.com  
www.neb.com



B0262S 002120915091

# B0262S

**6.0 ml Lambda Exonuclease Reaction Buffer (10X)**

**Store at -20°C Lot: 0021209 Exp: 9/15**

**Description:** New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

### **1X Lambda Exonuclease Reaction Buffer:**

67 mM Glycine-KOH  
2.5 mM MgCl<sub>2</sub>  
50 µg/ml BSA  
pH 9.4 @ 25°C

### **Quality Control Assays**

**16-Hour Incubation:** A 50 µl reaction containing this reaction buffer at a 1X concentration and 1 µg of HaeIII digested φX174 RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

**Endonuclease Activity:** Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reactions resulted in less than 5% conversion to RF II.

CERTIFICATE OF ANALYSIS