

Lambda Exonuclease Reaction Buffer



1-800-632-7799
info@neb.com
www.neb.com



B0262S 003130616061

B0262S

6.0 ml Lambda Exonuclease Reaction Buffer (10X)

Store at -20°C Lot: 0031306 Exp: 6/16

Description: New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X Lambda Exonuclease Reaction Buffer:

67 mM Glycine-KOH
2.5 mM MgCl₂
50 µg/ml BSA
pH 9.4 @ 25°C

Quality Control Assays

16-Hour Incubation: A 50 µl reaction containing this reaction buffer at a 1X concentration and 1 µg of HaeIII digested φX174 RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reactions resulted in less than 5% conversion to RF II.

CERTIFICATE OF ANALYSIS

Lambda Exonuclease Reaction Buffer



1-800-632-7799
info@neb.com
www.neb.com



B0262S 003130616061

B0262S

6.0 ml Lambda Exonuclease Reaction Buffer (10X)

Store at -20°C Lot: 0031306 Exp: 6/16

Description: New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X Lambda Exonuclease Reaction Buffer:

67 mM Glycine-KOH
2.5 mM MgCl₂
50 µg/ml BSA
pH 9.4 @ 25°C

Quality Control Assays

16-Hour Incubation: A 50 µl reaction containing this reaction buffer at a 1X concentration and 1 µg of HaeIII digested φX174 RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reactions resulted in less than 5% conversion to RF II.

CERTIFICATE OF ANALYSIS