

UDG Reaction Buffer



1-800-632-7799
info@neb.com
www.neb.com



B0280S 001130116011

B0280S

6.0 ml UDG Reaction Buffer (10X)

Store at -20°C Lot: 0011301 Exp: 1/16

Description: New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X UDG Reaction Buffer:

20 mM Tris-HCl
1 mM EDTA
1 mM dithiothreitol
(pH 8.0 @ 25°C)

Quality Control Assays

16-Hour Incubation: A 50 µl reaction containing this reaction buffer at a 1X concentration and 1 µg of HaeIII digested φX174 RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reactions resulted in less than 5% conversion to RF II.

CERTIFICATE OF ANALYSIS

UDG Reaction Buffer



1-800-632-7799
info@neb.com
www.neb.com



B0280S 001130116011

B0280S

6.0 ml UDG Reaction Buffer (10X)

Store at -20°C Lot: 0011301 Exp: 1/16

Description: New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X UDG Reaction Buffer:

20 mM Tris-HCl
1 mM EDTA
1 mM dithiothreitol
(pH 8.0 @ 25°C)

Quality Control Assays

16-Hour Incubation: A 50 µl reaction containing this reaction buffer at a 1X concentration and 1 µg of HaeIII digested φX174 RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reactions resulted in less than 5% conversion to RF II.

CERTIFICATE OF ANALYSIS