

NEBNext® dsDNA
Fragmentase®
Reaction Buffer



1-800-632-7799
info@neb.com
www.neb.com



B0348S 011160518051

B0348S

6 x 1.0 ml (6.0 ml) rxn buffer Lot: 0111605

Store at -20°C Exp: 5/18

Description: New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X NEBNext dsDNA Fragmentase Reaction Buffer:

20 mM Tris-HCl
10 mM MgCl₂
50 mM NaCl
0.15% Triton X-100
pH 7.5 @ 25°C

Quality Control Assay

16-Hour Incubation: A 50 µl reaction containing this reaction buffer at a 1X concentration and 1 µg of HaeIII digested φX174 RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reactions resulted in less than 5% conversion to RF II.

Protease Assay: Incubation of at least 1X dsDNA Fragmentase buffer with 0.2 nmol of a standardized mixture of proteins, for 20 hours at 37°C, resulted in no proteolytic activity detected by SDS-PAGE.

Phosphatase Assay: Incubation of 10 µl of at least 1X dsDNA Fragmentase Buffer in protein phosphatase assay buffer (1 M diethanolamine @ pH 9.8 and 0.5 mM MgCl₂) containing 2.5 mM *p*-nitrophenyl phosphate at 37°C for 4 hours yields no detectable *p*-nitrophenylene anion as determined by spectrophotometric analysis at 405 nm.



NEW ENGLAND BIOLABS®, FRAGMENTASE® and NEBNEXT® are registered trademarks of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

NEBNext® dsDNA
Fragmentase®
Reaction Buffer



1-800-632-7799
info@neb.com
www.neb.com



B0348S 011160518051

B0348S

6 x 1.0 ml (6.0 ml) rxn buffer Lot: 0111605

Store at -20°C Exp: 5/18

Description: New England Biolabs supplies a 10X reaction buffer with all of its enzymes. At a 1X concentration this reaction buffer assures optimal activity of the enzyme.

1X NEBNext dsDNA Fragmentase Reaction Buffer:

20 mM Tris-HCl
10 mM MgCl₂
50 mM NaCl
0.15% Triton X-100
pH 7.5 @ 25°C

Quality Control Assay

16-Hour Incubation: A 50 µl reaction containing this reaction buffer at a 1X concentration and 1 µg of HaeIII digested φX174 RF I DNA incubated for 16 hours resulted in no detectable non-specific nuclease degradation.

Endonuclease Activity: Incubation of this reaction buffer at a 1X concentration with 1 µg φX174 RF I DNA for 4 hours at 37°C in 50 µl reactions resulted in less than 5% conversion to RF II.

Protease Assay: Incubation of at least 1X dsDNA Fragmentase buffer with 0.2 nmol of a standardized mixture of proteins, for 20 hours at 37°C, resulted in no proteolytic activity detected by SDS-PAGE.

Phosphatase Assay: Incubation of 10 µl of at least 1X dsDNA Fragmentase Buffer in protein phosphatase assay buffer (1 M diethanolamine @ pH 9.8 and 0.5 mM MgCl₂) containing 2.5 mM *p*-nitrophenyl phosphate at 37°C for 4 hours yields no detectable *p*-nitrophenylene anion as determined by spectrophotometric analysis at 405 nm.



NEW ENGLAND BIOLABS®, FRAGMENTASE® and NEBNEXT® are registered trademarks of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS