

## NEBuffer 2.1



1-800-632-7799  
info@neb.com  
www.neb.com



B7202S 022170220021

# B7202S

5.0 ml Lot: 0221702  
Store at  $-20^{\circ}\text{C}$  Exp: 2/20

**Description:** New England Biolabs provides a color-coded 10X NEBuffer with each restriction endonuclease to ensure optimal (100%) activity. Most of our enzymes are supplied with one of four standard NEBuffers. Occasionally, an enzyme has specific buffer requirements not met by one of the four standard NEBuffers, in which case the enzyme is supplied with its own unique NEBuffer.

## NEBuffer 2.1



1-800-632-7799  
info@neb.com  
www.neb.com



B7202S 022170220021

# B7202S

5.0 ml Lot: 0221702  
Store at  $-20^{\circ}\text{C}$  Exp: 2/20

**Description:** New England Biolabs provides a color-coded 10X NEBuffer with each restriction endonuclease to ensure optimal (100%) activity. Most of our enzymes are supplied with one of four standard NEBuffers. Occasionally, an enzyme has specific buffer requirements not met by one of the four standard NEBuffers, in which case the enzyme is supplied with its own unique NEBuffer.

### 1X NEBuffer 2.1:

50 mM NaCl  
10 mM Tris-HCl  
10 mM MgCl<sub>2</sub>  
100 µg/ml BSA  
pH 7.9 @ 25°C  
Supplied as a 10X concentrated stock

### Quality Control

**pH range:** The pH of 10X NEBuffer 2.1 is between pH 7.8 and 8.0.

### 16-hour Non-specific Nuclease Activity Assay:

A 50 µl reaction in 1X NEBuffer 2.1 containing 1 µg of  $\phi$ X HaeIII digested DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**Endonuclease (nicking) Activity Assay:** A 50 µl reaction in 1X NEBuffer 2.1 containing 1 µg of supercoiled  $\phi$ X174 DNA incubated for 4 hours at 37°C results in < 10% conversion to the nicked form as determined by agarose gel electrophoresis.

### 1X NEBuffer 2.1:

50 mM NaCl  
10 mM Tris-HCl  
10 mM MgCl<sub>2</sub>  
100 µg/ml BSA  
pH 7.9 @ 25°C  
Supplied as a 10X concentrated stock

### Quality Control

**pH range:** The pH of 10X NEBuffer 2.1 is between pH 7.8 and 8.0.

### 16-hour Non-specific Nuclease Activity Assay:

A 50 µl reaction in 1X NEBuffer 2.1 containing 1 µg of  $\phi$ X HaeIII digested DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**Endonuclease (nicking) Activity Assay:** A 50 µl reaction in 1X NEBuffer 2.1 containing 1 µg of supercoiled  $\phi$ X174 DNA incubated for 4 hours at 37°C results in < 10% conversion to the nicked form as determined by agarose gel electrophoresis.

**Buffer Functional Assay:** A 50 µl reaction in 1X NEBuffer 2.1 containing 1 µg of  $\lambda$  DNA and 1 unit of either SphI or HindIII, incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.

**RNase Activity (Extended Digestion):** A 10 µl reaction in 1X NEBuffer 2.1 with 40 ng RNA transcript is incubated for 16 hours at 37°C. After incubation for 16 hours, no detectable degradation of the RNA is observed as determined by gel electrophoresis using fluorescent detection.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

**Buffer Functional Assay:** A 50 µl reaction in 1X NEBuffer 2.1 containing 1 µg of  $\lambda$  DNA and 1 unit of either SphI or HindIII, incubated for 1 hour at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.

**RNase Activity (Extended Digestion):** A 10 µl reaction in 1X NEBuffer 2.1 with 40 ng RNA transcript is incubated for 16 hours at 37°C. After incubation for 16 hours, no detectable degradation of the RNA is observed as determined by gel electrophoresis using fluorescent detection.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

CERTIFICATE OF ANALYSIS