

## Anti-M13 pIII Monoclonal Antibody



1-800-632-7799  
info@neb.com  
www.neb.com



E8033S 005121014101

# E8033S

**0.1 ml**      **Lot: 0051210**    **Exp: 10/14**  
**0.2 mg/ml**    **Store at -20°C**

**Description:** Anti-M13 pIII Monoclonal Antibody (mouse isotype IgG2a) is derived from the A23 hybridoma produced by the fusion of mouse myeloma cells and splenocytes from BALB/c mice immunized with the C-terminal half of M13 coat protein III (residues 259–406). Monoclonal anti-M13 pIII recognizes nonreduced and denatured reduced forms of wild type pIII or pIII fusions in immunoblotting and ELISA.

## Anti-M13 pIII Monoclonal Antibody



1-800-632-7799  
info@neb.com  
www.neb.com



E8033S 005121014101

# E8033S

**0.1 ml**      **Lot: 0051210**    **Exp: 10/14**  
**0.2 mg/ml**    **Store at -20°C**

**Description:** Anti-M13 pIII Monoclonal Antibody (mouse isotype IgG2a) is derived from the A23 hybridoma produced by the fusion of mouse myeloma cells and splenocytes from BALB/c mice immunized with the C-terminal half of M13 coat protein III (residues 259–406). Monoclonal anti-M13 pIII recognizes nonreduced and denatured reduced forms of wild type pIII or pIII fusions in immunoblotting and ELISA.

**Source:** Tissue culture supernatant from murine monoclonal cell line A23.

Supplied in: 10 mM HEPES (pH 7.5),  
150 mM NaCl and 50% glycerol.

**Applications:** Recombinant protein domains such as antibody fragments can be fused to M13 coat protein pIII and displayed on phage using phagemid-based systems (1), rather than the phage-based system utilized in the Ph.D. libraries. Anti-M13 pIII Monoclonal Antibody can be used to determine fusion protein expression levels both in *E. coli* and on the M13 virion by immunoblot analysis.

**Suggested Working Dilution:** 1/1,000

**Performance:** A dilution of 1/1,000 detected as few as  $10^7$  M13 CFU electrophoresed and transferred onto Immobolin PVDF membrane in an immunoblot assay.

**Source:** Tissue culture supernatant from murine monoclonal cell line A23.

Supplied in: 10 mM HEPES (pH 7.5),  
150 mM NaCl and 50% glycerol.

**Applications:** Recombinant protein domains such as antibody fragments can be fused to M13 coat protein pIII and displayed on phage using phagemid-based systems (1), rather than the phage-based system utilized in the Ph.D. libraries. Anti-M13 pIII Monoclonal Antibody can be used to determine fusion protein expression levels both in *E. coli* and on the M13 virion by immunoblot analysis.

**Suggested Working Dilution:** 1/1,000

**Performance:** A dilution of 1/1,000 detected as few as  $10^7$  M13 CFU electrophoresed and transferred onto Immobolin PVDF membrane in an immunoblot assay.

**Note:** ELISA against whole phage using this antibody is not recommended since the epitope lies in a region that is not accessible on intact M13 phage virions.

### Reference:

1. Barbas, C.F. et al. (2001). *Phage Display: A Laboratory Manual*, (pp. 2.1–2.19). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.

**Note:** ELISA against whole phage using this antibody is not recommended since the epitope lies in a region that is not accessible on intact M13 phage virions.

### Reference:

1. Barbas, C.F. et al. (2001). *Phage Display: A Laboratory Manual*, (pp. 2.1–2.19). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.

CERTIFICATE OF ANALYSIS

CERTIFICATE OF ANALYSIS