

Anti-M13 pIII Monoclonal Antibody



1-800-632-7799
info@neb.com
www.neb.com



E8033S 006151217121

E8033S

0.1 ml **Lot: 0061512** **Exp: 12/17**
0.2 mg/ml **Store at -20°C**

Description: Anti-M13 pIII Monoclonal Antibody (mouse isotype IgG2a) is derived from the A23 hybridoma produced by the fusion of mouse myeloma cells and splenocytes from BALB/c mice immunized with the C-terminal half of M13 coat protein III (residues 259–406). Monoclonal anti-M13 pIII recognizes nonreduced and denatured reduced forms of wild type pIII or pIII fusions in immunoblotting and ELISA.

LABORATORY REAGENT – FOR RESEARCH USE ONLY

Source: Tissue culture supernatant from murine monoclonal cell line A23.

Supplied in: 10 mM HEPES (pH 7.5), 150 mM NaCl and 50% glycerol.

Applications: Recombinant protein domains such as antibody fragments can be fused to M13 coat protein pIII and displayed on phage using phagemid-based systems (1), rather than the phage-based system utilized in the Ph.D. libraries. Anti-M13 pIII Monoclonal Antibody can be used to determine fusion protein expression levels both in *E. coli* and on the M13 virion by immunoblot analysis.

Suggested Working Dilution: 1/1,000

Performance: A dilution of 1/1,000 detected as few as 10⁷ M13 CFU electrophoresed and transferred onto Immobolin PVDF membrane in an immunoblot assay.

Note: ELISA against whole phage using this antibody is not recommended since the epitope lies in a region that is not accessible on intact M13 phage virions.

Reference:

1. Barbas, C.F. et al. (2001). *Phage Display: A Laboratory Manual*, (pp. 2.1–2.19). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

Anti-M13 pIII Monoclonal Antibody



1-800-632-7799
info@neb.com
www.neb.com



E8033S 006151217121

E8033S

0.1 ml **Lot: 0061512** **Exp: 12/17**
0.2 mg/ml **Store at -20°C**

Description: Anti-M13 pIII Monoclonal Antibody (mouse isotype IgG2a) is derived from the A23 hybridoma produced by the fusion of mouse myeloma cells and splenocytes from BALB/c mice immunized with the C-terminal half of M13 coat protein III (residues 259–406). Monoclonal anti-M13 pIII recognizes nonreduced and denatured reduced forms of wild type pIII or pIII fusions in immunoblotting and ELISA.

LABORATORY REAGENT – FOR RESEARCH USE ONLY

Source: Tissue culture supernatant from murine monoclonal cell line A23.

Supplied in: 10 mM HEPES (pH 7.5), 150 mM NaCl and 50% glycerol.

Applications: Recombinant protein domains such as antibody fragments can be fused to M13 coat protein pIII and displayed on phage using phagemid-based systems (1), rather than the phage-based system utilized in the Ph.D. libraries. Anti-M13 pIII Monoclonal Antibody can be used to determine fusion protein expression levels both in *E. coli* and on the M13 virion by immunoblot analysis.

Suggested Working Dilution: 1/1,000

Performance: A dilution of 1/1,000 detected as few as 10⁷ M13 CFU electrophoresed and transferred onto Immobolin PVDF membrane in an immunoblot assay.

Note: ELISA against whole phage using this antibody is not recommended since the epitope lies in a region that is not accessible on intact M13 phage virions.

Reference:

1. Barbas, C.F. et al. (2001). *Phage Display: A Laboratory Manual*, (pp. 2.1–2.19). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS