

T4 RNA Ligase 1 (ssRNA Ligase)



1-800-632-7799
info@neb.com
www.neb.com



M0204S 060160418041

M0204S



1,000 units **10,000 U/ml** **Lot: 0601604**

RECOMBINANT **Store at -20°C** **Exp: 4/18**

Description: T4 RNA Ligase 1 catalyzes the ligation of a 5' phosphoryl-terminated nucleic acid donor to a 3' hydroxyl-terminated nucleic acceptor through the formation of a 3' → 5' phosphodiester bond, with hydrolysis of ATP to AMP and PP_i. Substrates include single-stranded RNA and DNA as well as dinucleoside pyrophosphates (1).

Source: An *E. coli* strain that carries the T4 RNA Ligase 1 gene

Applications:

- Ligation of ss-RNA and DNA
- Labeling of 3'-termini of RNA with 5'-[³²P] pCp (3)
- Inter- and intramolecular joining of RNA and DNA molecules (4,5)
- Synthesis of single-stranded oligodeoxyribonucleotides (6)
- Incorporation of unnatural amino acids into proteins (7)

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 1 mM DTT and 50% glycerol.

Reagents Supplied with Enzyme:

10X T4 RNA Ligase Reaction Buffer, 10 mM ATP and 50% PEG 8000.

Reaction Conditions: 1X T4 RNA Ligase Reaction Buffer, supplemented with 1 mM ATP. Incubate at 37°C.

1X T4 RNA Ligase Reaction Buffer:

50 mM Tris-HCl
10 mM MgCl₂
1 mM DTT
pH 7.5 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to convert 1 nanomole of 5'-[³²P] rA₁₆ into a phosphatase-resistant form in 30 minutes at 37°C

Unit Assay Conditions: 1X T4 RNA Ligase reaction buffer, supplemented with 1 mM ATP, is mixed with the RNA substrate (10 μM of 5'-[³²P]rA₁₆) and varying amounts of enzyme. Incubation is at 37°C for 15 minutes (8).

Heat Inactivation: 65°C for 15 minutes or boiling for 2 minutes.

Quality Control Assays

RNase Assay: Incubation of a 10 μl reaction containing 20 units of T4 RNA Ligase 1 with 40 ng of RNA transcript for 2 hours at 37°C resulted in no detectable degradation of the RNA as determined by gel electrophoresis.

DNA Exonuclease Activity: Incubation of 20 units of T4 RNA Ligase 1 with 1 μg of mixed single and double-stranded sonicated ³H DNA (10⁵ cpm/μg) in 50 μl T4 RNA Ligase Reaction Buffer for 4 hours at 37°C released < 0.1% of the activity.

DNA Endonuclease Activity: Incubation of 20 units of T4 RNA Ligase 1 with 1 μg φX174 RF I DNA in 50 μl T4 RNA Ligase Reaction Buffer for 4 hours at 37°C resulted in no detectable degradation of DNA as determined by agarose gel electrophoresis.

Notes on Use: Addition of DMSO to 10% (v/v) is required for pCp ligation (3).

(see other side)

CERTIFICATE OF ANALYSIS

T4 RNA Ligase 1 (ssRNA Ligase)



1-800-632-7799
info@neb.com
www.neb.com



M0204S 060160418041

M0204S



1,000 units **10,000 U/ml** **Lot: 0601604**

RECOMBINANT **Store at -20°C** **Exp: 4/18**

Description: T4 RNA Ligase 1 catalyzes the ligation of a 5' phosphoryl-terminated nucleic acid donor to a 3' hydroxyl-terminated nucleic acceptor through the formation of a 3' → 5' phosphodiester bond, with hydrolysis of ATP to AMP and PP_i. Substrates include single-stranded RNA and DNA as well as dinucleoside pyrophosphates (1).

Source: An *E. coli* strain that carries the T4 RNA Ligase 1 gene

Applications:

- Ligation of ss-RNA and DNA
- Labeling of 3'-termini of RNA with 5'-[³²P] pCp (3)
- Inter- and intramolecular joining of RNA and DNA molecules (4,5)
- Synthesis of single-stranded oligodeoxyribonucleotides (6)
- Incorporation of unnatural amino acids into proteins (7)

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 1 mM DTT and 50% glycerol.

Reagents Supplied with Enzyme:

10X T4 RNA Ligase Reaction Buffer, 10 mM ATP and 50% PEG 8000.

Reaction Conditions: 1X T4 RNA Ligase Reaction Buffer, supplemented with 1 mM ATP. Incubate at 37°C.

1X T4 RNA Ligase Reaction Buffer:

50 mM Tris-HCl
10 mM MgCl₂
1 mM DTT
pH 7.5 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to convert 1 nanomole of 5'-[³²P] rA₁₆ into a phosphatase-resistant form in 30 minutes at 37°C

Unit Assay Conditions: 1X T4 RNA Ligase reaction buffer, supplemented with 1 mM ATP, is mixed with the RNA substrate (10 μM of 5'-[³²P]rA₁₆) and varying amounts of enzyme. Incubation is at 37°C for 15 minutes (8).

Heat Inactivation: 65°C for 15 minutes or boiling for 2 minutes.

Quality Control Assays

RNase Assay: Incubation of a 10 μl reaction containing 20 units of T4 RNA Ligase 1 with 40 ng of RNA transcript for 2 hours at 37°C resulted in no detectable degradation of the RNA as determined by gel electrophoresis.

DNA Exonuclease Activity: Incubation of 20 units of T4 RNA Ligase 1 with 1 μg of mixed single and double-stranded sonicated ³H DNA (10⁵ cpm/μg) in 50 μl T4 RNA Ligase Reaction Buffer for 4 hours at 37°C released < 0.1% of the activity.

DNA Endonuclease Activity: Incubation of 20 units of T4 RNA Ligase 1 with 1 μg φX174 RF I DNA in 50 μl T4 RNA Ligase Reaction Buffer for 4 hours at 37°C resulted in no detectable degradation of DNA as determined by agarose gel electrophoresis.

Notes on Use: Addition of DMSO to 10% (v/v) is required for pCp ligation (3).

(see other side)

CERTIFICATE OF ANALYSIS

References:

1. England, T., Gumport, R. and Uhlenbeck, O. (1977) *Proc. Natl. Acad. Sci. USA* 74, 4839–4842.
2. Rand, K.N. and Gait, M.J. (1984) *EMBO J.* 3, 397–402.
3. England, T. and Uhlenbeck, O. (1978) *Nature* 275, 560–562.
4. Romaniuk, P. and Uhlenbeck, O. (1983). In R. Wu, L. Grossman and K. Moldave (Eds.), *Methods in Enzymology* Vol. 100, (pp.52–56). New York: Academic Press.
5. Moore, M.J. and Sharp, P.A. (1992) *Science* 256, 992–997.
6. Tessier, D.C., Brousseau, R. and Vernet, T. (1986) *Anal. Biochem.* 158, 171–178.
7. Noren, C.J. et al. (1989) *Science* 244, 182–188.
8. Silber, R., Malathi, B.G. and Hurwitz, J. (1972). *Proc. Natl. Acad. Sci. USA* 69, 3009–3013.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

References:

1. England, T., Gumport, R. and Uhlenbeck, O. (1977) *Proc. Natl. Acad. Sci. USA* 74, 4839–4842.
2. Rand, K.N. and Gait, M.J. (1984) *EMBO J.* 3, 397–402.
3. England, T. and Uhlenbeck, O. (1978) *Nature* 275, 560–562.
4. Romaniuk, P. and Uhlenbeck, O. (1983). In R. Wu, L. Grossman and K. Moldave (Eds.), *Methods in Enzymology* Vol. 100, (pp.52–56). New York: Academic Press.
5. Moore, M.J. and Sharp, P.A. (1992) *Science* 256, 992–997.
6. Tessier, D.C., Brousseau, R. and Vernet, T. (1986) *Anal. Biochem.* 158, 171–178.
7. Noren, C.J. et al. (1989) *Science* 244, 182–188.
8. Silber, R., Malathi, B.G. and Hurwitz, J. (1972). *Proc. Natl. Acad. Sci. USA* 69, 3009–3013.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.