

E. coli DNA Ligase



1-800-632-7799
info@neb.com
www.neb.com



M0205S 081160918091

M0205S



200 units Lot: **0811609** Exp: **9/18**
10,000 U/ml Store at **-20°C**

Description: Catalyzes the formation of a phosphodiester bond between juxtaposed 5'-phosphate and 3'-hydroxyl termini in duplex DNA containing cohesive ends.

Source: Purified from *E. coli* strain containing a cloned *E. coli* DNA Ligase gene.

E. coli DNA Ligase



1-800-632-7799
info@neb.com
www.neb.com



M0205S 081160918091

M0205S



200 units Lot: **0811609** Exp: **9/18**
10,000 U/ml Store at **-20°C**

Description: Catalyzes the formation of a phosphodiester bond between juxtaposed 5'-phosphate and 3'-hydroxyl termini in duplex DNA containing cohesive ends.

Source: Purified from *E. coli* strain containing a cloned *E. coli* DNA Ligase gene.

Applications:

- Okayama and Berg cDNA cloning (3)

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X *E. coli* DNA Ligase Buffer.

Reaction Conditions: Incubate DNA at a final concentration of 0.12 µM in 20–50 µl of 1X *E. coli* DNA Ligase Buffer at 16°C overnight.

1X *E. coli* DNA Ligase Buffer:

30 mM Tris-HCl
4 mM MgCl₂
1 mM DTT
26 µM NAD⁺
50 µg/ml BSA
pH 8.0 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to give 50% ligation of Hind III fragments of λ DNA (5' DNA termini concentration of 0.12 µM, 300 µg/ml) in a total reaction volume of 20 µl in 30 minutes at 16°C.

Applications:

- Okayama and Berg cDNA cloning (3)

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM DTT, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X *E. coli* DNA Ligase Buffer.

Reaction Conditions: Incubate DNA at a final concentration of 0.12 µM in 20–50 µl of 1X *E. coli* DNA Ligase Buffer at 16°C overnight.

1X *E. coli* DNA Ligase Buffer:

30 mM Tris-HCl
4 mM MgCl₂
1 mM DTT
26 µM NAD⁺
50 µg/ml BSA
pH 8.0 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to give 50% ligation of Hind III fragments of λ DNA (5' DNA termini concentration of 0.12 µM, 300 µg/ml) in a total reaction volume of 20 µl in 30 minutes at 16°C.

Unit Assay Conditions: 30 mM Tris-HCl (pH 8.0), 4 mM MgCl₂, 1 mM dithiothreitol, 26 µM NAD⁺, 50 µg/ml bovine serum albumin and Hind III fragments of λ DNA (300 µg/ml).

Specific Activity: ~ 6,000 units/mg.

Heat Inactivation: 65°C for 20 minutes.

Quality Control Assays

Exonuclease Activity: Incubation of 20 units of enzyme for 4 hours at 37°C in 50 µl assay buffer containing 1 µg sonicated *E. coli* ³H DNA (10⁵ cpm/µg) resulted in < 0.1% acid soluble counts.

16 Hour-Incubation: Incubation of 20 units of *E. coli* DNA Ligase with 1 µg of λ DNA (HindIII digest) for 16 hours in 50 µl of 1X NEBuffer 3 at 37°C resulted in a normal and sharp banding pattern on agarose gels.

Notes on Use: Requires NAD⁺ (nicotinamide adenine dinucleotide) as a cofactor, in contrast to other ligases which use rATP.

Unit Assay Conditions: 30 mM Tris-HCl (pH 8.0), 4 mM MgCl₂, 1 mM dithiothreitol, 26 µM NAD⁺, 50 µg/ml bovine serum albumin and Hind III fragments of λ DNA (300 µg/ml).

Specific Activity: ~ 6,000 units/mg.

Heat Inactivation: 65°C for 20 minutes.

Quality Control Assays

Exonuclease Activity: Incubation of 20 units of enzyme for 4 hours at 37°C in 50 µl assay buffer containing 1 µg sonicated *E. coli* ³H DNA (10⁵ cpm/µg) resulted in < 0.1% acid soluble counts.

16 Hour-Incubation: Incubation of 20 units of *E. coli* DNA Ligase with 1 µg of λ DNA (HindIII digest) for 16 hours in 50 µl of 1X NEBuffer 3 at 37°C resulted in a normal and sharp banding pattern on agarose gels.

Notes on Use: Requires NAD⁺ (nicotinamide adenine dinucleotide) as a cofactor, in contrast to other ligases which use rATP.

Ligation of blunt-ended fragments is extremely inefficient. For ligation of blunt-ended fragments use T4 DNA Ligase.

Does not ligate RNA to DNA (4).

This enzyme ligates only DNA fragments with cohesive termini.

References:

1. Panasenko, S. M. et al. (1977) *Science* 196, 188–189.
2. Panasenko, S. M. et al. (1978) *J. Biol. Chem.* 253, 4590–4592.
3. Okayama, H. and Berg, P. (1982) *Mol. Cell. Biol.* 2, 161–170.
4. Higgins, N. P. and Cozzarelli, N. R. (1979) *Methods Enzymol.* 68, 50–71.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

Ligation of blunt-ended fragments is extremely inefficient. For ligation of blunt-ended fragments use T4 DNA Ligase.

Does not ligate RNA to DNA (4).

This enzyme ligates only DNA fragments with cohesive termini.

References:

1. Panasenko, S. M. et al. (1977) *Science* 196, 188–189.
2. Panasenko, S. M. et al. (1978) *J. Biol. Chem.* 253, 4590–4592.
3. Okayama, H. and Berg, P. (1982) *Mol. Cell. Biol.* 2, 161–170.
4. Higgins, N. P. and Cozzarelli, N. R. (1979) *Methods Enzymol.* 68, 50–71.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS