

SP6 RNA Polymerase



1-800-632-7799
info@neb.com
www.neb.com



M0207S 015160418041

M0207S



2,000 units 20,000 U/ml Lot: 0151604
RECOMBINANT Store at -20°C Exp: 4/18

Description: Bacteriophage SP6 RNA Polymerase is a DNA-dependent RNA polymerase that is highly specific for the SP6 phage promoter. The 98.5 KD polymerase catalyzes *in vitro* RNA synthesis from a cloned DNA template under the SP6 promoter. RNA synthesized using the SP6 RNA Polymerase is suitable for many applications in research and biotechnology.

Source: An *E. coli* strain that carries the cloned gene for SP6 RNA Polymerase from *Salmonella typhimurium* LT2Z.

Applications:

- Radiolabeled RNA probe preparation
- Non-isotopic RNA labeling
- Preparation of RNA vaccines
- Guide RNA for gene targeting
- mRNA for *in vitro* translation and micro injection
- RNA structure, processing and catalysis studies
- RNA amplification
- Anti-sense RNA for gene expression experiment

Supplied in: 100 mM NaCl, 50 mM Tris-HCl (pH 7.9), 1 mM EDTA, 20 mM 2-mercaptoethanol, 0.1% Triton X-100 and 50% glycerol.

Reagents Supplied with Enzyme:

10X RNAPol Reaction Buffer.

Reaction Conditions: 1X RNAPol Reaction Buffer, supplemented with 0.5 mM each ATP, UTP, GTP, CTP and DNA template containing the SP6 RNA Polymerase promoter. Incubate at 37°C.

Applications:

- Radiolabeled RNA probe preparation
- Non-isotopic RNA labeling
- Preparation of RNA vaccines
- Guide RNA for gene targeting
- mRNA for *in vitro* translation and micro injection
- RNA structure, processing and catalysis studies
- RNA amplification
- Anti-sense RNA for gene expression experiment

Supplied in: 100 mM NaCl, 50 mM Tris-HCl (pH 7.9), 1 mM EDTA, 20 mM 2-mercaptoethanol, 0.1% Triton X-100 and 50% glycerol.

Reagents Supplied with Enzyme:

10X RNAPol Reaction Buffer.

Reaction Conditions: 1X RNAPol Reaction Buffer, supplemented with 0.5 mM each ATP, UTP, GTP, CTP and DNA template containing the SP6 RNA Polymerase promoter. Incubate at 37°C.

1X RNAPol Reaction Buffer:

40 mM Tris-HCl
6 mM MgCl₂
2 mM spermidine
1 mM dithiothreitol
(pH 7.9 @ 25°C)

Unit Definition: One unit is defined as the amount of enzyme required to incorporate 1 nmol ATP into an acid-insoluble material in 1 hour at 37°C.

Unit Assay Conditions: 1X RNAPol Reaction Buffer, supplemented with 0.5 mM each ATP, UTP, GTP, CTP and 1 µg DNA containing the SP6 promoter in 50 µl.

Protocol for Standard RNA Synthesis: Assemble the reaction at room temperature in the following order.

COMPONENTS	AMOUNT	CONCENTRATION
H ₂ O	X µl	
10X Reaction Buffer	2 µl	
NTP	X µl	0.5 mM each
Template DNA	X µl	0.2–1 µg
RNase Inhibitor (optional)	0.5 µl	1 U/µl final
Fresh DTT (optional)	X µl	5 mM final
SP6 RNA Pol	2 µl	
Total Volume	20 µl	

1X RNAPol Reaction Buffer:

40 mM Tris-HCl
6 mM MgCl₂
2 mM spermidine
1 mM dithiothreitol
(pH 7.9 @ 25°C)

Unit Definition: One unit is defined as the amount of enzyme required to incorporate 1 nmol ATP into an acid-insoluble material in 1 hour at 37°C.

Unit Assay Conditions: 1X RNAPol Reaction Buffer, supplemented with 0.5 mM each ATP, UTP, GTP, CTP and 1 µg DNA containing the SP6 promoter in 50 µl.

Protocol for Standard RNA Synthesis: Assemble the reaction at room temperature in the following order.

COMPONENTS	AMOUNT	CONCENTRATION
H ₂ O	X µl	
10X Reaction Buffer	2 µl	
NTP	X µl	0.5 mM each
Template DNA	X µl	0.2–1 µg
RNase Inhibitor (optional)	0.5 µl	1 U/µl final
Fresh DTT (optional)	X µl	5 mM final
SP6 RNA Pol	2 µl	
Total Volume	20 µl	

Incubate at 37°C for 1 hour. For shorter (< 300 nt) transcripts incubate at 37°C for 2–16 hours.

Notes on use:

For radio labeled high specific activity RNA probes, the concentration of the radioactive nucleotide should be limited to 6 µM.

To protect RNA against ribonuclease, RNase inhibitor (NEB #M0314 or #M0307) should be added to a final concentration of 1 U/µl.

SP6 RNA Polymerase is extremely sensitive to salt inhibition. The overall salt concentration should not exceed 50 mM.

SP6 RNA Polymerase is slightly more active at 40°C than at 37°C. Incubation at 40°C may be considered for RNA transcript containing strong secondary structures.

Quality Control Assays

Endonuclease Activity (Nicking): A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of supercoiled ϕX174 DNA and a minimum of 100 units of SP6 RNA Polymerase incubated for 4 hours at

(see other side)

CERTIFICATE OF ANALYSIS

SP6 RNA Polymerase



1-800-632-7799
info@neb.com
www.neb.com



M0207S 015160418041

M0207S



2,000 units 20,000 U/ml Lot: 0151604
RECOMBINANT Store at -20°C Exp: 4/18

Description: Bacteriophage SP6 RNA Polymerase is a DNA-dependent RNA polymerase that is highly specific for the SP6 phage promoter. The 98.5 KD polymerase catalyzes *in vitro* RNA synthesis from a cloned DNA template under the SP6 promoter. RNA synthesized using the SP6 RNA Polymerase is suitable for many applications in research and biotechnology.

Source: An *E. coli* strain that carries the cloned gene for SP6 RNA Polymerase from *Salmonella typhimurium* LT2Z.

Incubate at 37°C for 1 hour. For shorter (< 300 nt) transcripts incubate at 37°C for 2–16 hours.

Notes on use:

For radio labeled high specific activity RNA probes, the concentration of the radioactive nucleotide should be limited to 6 µM.

To protect RNA against ribonuclease, RNase inhibitor (NEB #M0314 or #M0307) should be added to a final concentration of 1 U/µl.

SP6 RNA Polymerase is extremely sensitive to salt inhibition. The overall salt concentration should not exceed 50 mM.

SP6 RNA Polymerase is slightly more active at 40°C than at 37°C. Incubation at 40°C may be considered for RNA transcript containing strong secondary structures.

Quality Control Assays

Endonuclease Activity (Nicking): A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of supercoiled ϕX174 DNA and a minimum of 100 units of SP6 RNA Polymerase incubated for 4 hours at

(see other side)

CERTIFICATE OF ANALYSIS

37°C results in < 10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release):

A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 100 units of SP6 RNA Polymerase incubated for 4 hours at 37°C releases < 0.1% of the total radioactivity.

Non-Specific DNase Activity (16 Hour): A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of SP6 RNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Promoter Specificity: A 50 µl reaction in RNAPol Reaction Buffer in the presence of 2 mM NTPs containing 1 µg of Lambda DNA as a template and a minimum of 100 units of SP6 RNA Polymerase incubated for 1 hour at 37°C results in < 1.5% of the amount of product incorporated as compared to a control reaction using SP6 DNA as a template.

37°C results in < 10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release):

A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 100 units of SP6 RNA Polymerase incubated for 4 hours at 37°C releases < 0.1% of the total radioactivity.

Non-Specific DNase Activity (16 Hour): A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of SP6 RNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Promoter Specificity: A 50 µl reaction in RNAPol Reaction Buffer in the presence of 2 mM NTPs containing 1 µg of Lambda DNA as a template and a minimum of 100 units of SP6 RNA Polymerase incubated for 1 hour at 37°C results in < 1.5% of the amount of product incorporated as compared to a control reaction using SP6 DNA as a template.

Protein Purity Assay (SDS-PAGE): SP6 RNA Polymerase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

RNase Activity (Extended Digestion): A 10 µl reaction in RNAPol Reaction Buffer containing 40 ng of a 300 base single-stranded RNA and a minimum of 20 units of SP6 RNA Polymerase is incubated at 37°C. After incubation for 4 hours, > 90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

Companion Products:

RNA Loading Dye (2X)
#B0363S 4 x 1 ml

RNase Inhibitor, Human Placenta
#M0307S 2,000 units
#M0307L 10,000 units

RNase Inhibitor, Murine
#M0314S 3,000 units
#M0314L 15,000 units

Protein Purity Assay (SDS-PAGE): SP6 RNA Polymerase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

RNase Activity (Extended Digestion): A 10 µl reaction in RNAPol Reaction Buffer containing 40 ng of a 300 base single-stranded RNA and a minimum of 20 units of SP6 RNA Polymerase is incubated at 37°C. After incubation for 4 hours, > 90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

Companion Products:

RNA Loading Dye (2X)
#B0363S 4 x 1 ml

RNase Inhibitor, Human Placenta
#M0307S 2,000 units
#M0307L 10,000 units

RNase Inhibitor, Murine
#M0314S 3,000 units
#M0314L 15,000 units

Pyrophosphatase, Inorganic (*E. coli*)
#M0361S 10 units
#M0361L 50 units

Low Range ssRNA Ladder
#N0364S 25 µg

ssRNA Ladder
#N0362S 25 µg

RNase Contamination Assay Kit
#E3320S 50 reactions

Vaccinia Capping System
#M2080S 400 units

mRNA Cap 2'-O-Methyltransferase
#M0366S 2,000 units

E. coli Poly(A) Polymerase
#M0276S 100 units
#M0276L 500 units

Ribonucleotide Solution Mix
#N0446S 10 µmol of each
#N0446L 50 µmol of each

Pyrophosphatase, Inorganic (*E. coli*)
#M0361S 10 units
#M0361L 50 units

Low Range ssRNA Ladder
#N0364S 25 µg

ssRNA Ladder
#N0362S 25 µg

RNase Contamination Assay Kit
#E3320S 50 reactions

Vaccinia Capping System
#M2080S 400 units

mRNA Cap 2'-O-Methyltransferase
#M0366S 2,000 units

E. coli Poly(A) Polymerase
#M0276S 100 units
#M0276L 500 units

Ribonucleotide Solution Mix
#N0446S 10 µmol of each
#N0446L 50 µmol of each

Ribonucleotide Solution Set
#N0450S 10 µmol each
#N0450L 50 µmol each

T7 RNA Polymerase
#M0251S 5,000 units
#M0251L 25,000 units



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

Ribonucleotide Solution Set
#N0450S 10 µmol each
#N0450L 50 µmol each

T7 RNA Polymerase
#M0251S 5,000 units
#M0251L 25,000 units



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.