

Nuclease BAL-31



1-800-632-7799
info@neb.com
www.neb.com



M0213S 040151017101

M0213S



50 units **Lot: 0401510** **Exp: 10/17**
1,000 U/ml **Store at -20°C**

Description: BAL-31 exonuclease degrades both 3' and 5' termini of duplex DNA without generating internal scissions. The enzyme is also a highly specific single-stranded endonuclease which cleaves at nicks, gaps and single-stranded regions of duplex DNA and RNA (1,2).

Source: Purified from the culture medium of *Alteromonas espejiana* BAL-31. Contains a mixture of "fast" and "slow" species of the enzyme (3).

Applications:

- Progressive shortening of double-stranded DNA fragments at both termini (4)
- Restriction site mapping (2).

Supplied in: 50 mM NaCl, 10 mM Tris-HCl (pH 8.0), 1.5 mM CaCl₂, 1.5 mM MgCl₂, 0.25 mM EDTA, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

2X Nuclease BAL-31 Reaction Buffer.

Reaction Conditions: 1X Nuclease BAL-31 Reaction Buffer. **Incubate at 30°C.**

1X Nuclease BAL-31 Reaction Buffer:

600 mM NaCl
12 mM CaCl₂
12 mM MgCl₂
20 mM Tris-HCl
1 mM EDTA
pH 8.0 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to remove 200 base pairs from each end of linearized double-stranded φX174 DNA (40 µg/ml) in 50 µl of 1X Nuclease BAL-31 Reaction Buffer in 10 minutes at 30°C.

Applications:

- Progressive shortening of double-stranded DNA fragments at both termini (4)
- Restriction site mapping (2).

Supplied in: 50 mM NaCl, 10 mM Tris-HCl (pH 8.0), 1.5 mM CaCl₂, 1.5 mM MgCl₂, 0.25 mM EDTA, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

2X Nuclease BAL-31 Reaction Buffer.

Reaction Conditions: 1X Nuclease BAL-31 Reaction Buffer. **Incubate at 30°C.**

1X Nuclease BAL-31 Reaction Buffer:

600 mM NaCl
12 mM CaCl₂
12 mM MgCl₂
20 mM Tris-HCl
1 mM EDTA
pH 8.0 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to remove 200 base pairs from each end of linearized double-stranded φX174 DNA (40 µg/ml) in 50 µl of 1X Nuclease BAL-31 Reaction Buffer in 10 minutes at 30°C.

Heat Inactivation: Heat inactivated by incubation at 65°C for 10 minutes in the presence of 20 mM EGTA, a specific chelator of the essential cofactor Ca²⁺. This treatment does not affect the Mg²⁺ concentration.

Quality Control Assays

Ligation Activity: Incubation of 50 µl reaction buffer containing 30 units of enzyme and 25 µg HaeIII fragments of φX174 RF I DNA for 5 minutes at 30°C allowed the ligation of approximately 75% of the fragments during subsequent incubation with T4 DNA Ligase (5' termini concentration-20 µM).

Double-Stranded Endonuclease Activity:

Incubation of 60 units of Nuclease BAL-31 with 65 µg λ DNA for 10 minutes at 30°C in 100 µl reaction buffer (rendering 50% of the DNA acid-soluble) resulted in no detectable endonuclease activity. This is judged by the integrity of the internal λ DNA fragments produced by subsequent digestion with Hind III endonuclease.

Heat Inactivation: Heat inactivated by incubation at 65°C for 10 minutes in the presence of 20 mM EGTA, a specific chelator of the essential cofactor Ca²⁺. This treatment does not affect the Mg²⁺ concentration.

Quality Control Assays

Ligation Activity: Incubation of 50 µl reaction buffer containing 30 units of enzyme and 25 µg HaeIII fragments of φX174 RF I DNA for 5 minutes at 30°C allowed the ligation of approximately 75% of the fragments during subsequent incubation with T4 DNA Ligase (5' termini concentration-20 µM).

Double-Stranded Endonuclease Activity:

Incubation of 60 units of Nuclease BAL-31 with 65 µg λ DNA for 10 minutes at 30°C in 100 µl reaction buffer (rendering 50% of the DNA acid-soluble) resulted in no detectable endonuclease activity. This is judged by the integrity of the internal λ DNA fragments produced by subsequent digestion with Hind III endonuclease.

Notes On Use: Duplex products of the exonuclease are a mixture of blunt and staggered ends. This mixture can be cloned directly, although maximal ligation efficiency requires repairing the staggered ends with a suitable DNA polymerase.

If necessary, the enzyme may be diluted in reaction buffer prior to use.

Activity is linear with enzyme concentration.

References:

1. Gray, H. B. et al. (1975) *Nucleic Acids Res.* 2, 1459-1492.
2. Legerski, R. J. et al. (1978) *Nucleic Acids Res.* 5, 1445-1463.
3. Wei, C. -F. et al. (1983) *J. Biol. Chem.* 258, 13506-13512.
4. Sambrook, J., Fritsch, E. F. and Maniatis, T. (1989). *Molecular Cloning: A Laboratory Manual*, (2nd ed.), (pp. 5.73-5.75). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.

CERTIFICATE OF ANALYSIS

Nuclease BAL-31



1-800-632-7799
info@neb.com
www.neb.com



M0213S 040151017101

M0213S



50 units **Lot: 0401510** **Exp: 10/17**
1,000 U/ml **Store at -20°C**

Description: BAL-31 exonuclease degrades both 3' and 5' termini of duplex DNA without generating internal scissions. The enzyme is also a highly specific single-stranded endonuclease which cleaves at nicks, gaps and single-stranded regions of duplex DNA and RNA (1,2).

Source: Purified from the culture medium of *Alteromonas espejiana* BAL-31. Contains a mixture of "fast" and "slow" species of the enzyme (3).

Notes On Use: Duplex products of the exonuclease are a mixture of blunt and staggered ends. This mixture can be cloned directly, although maximal ligation efficiency requires repairing the staggered ends with a suitable DNA polymerase.

If necessary, the enzyme may be diluted in reaction buffer prior to use.

Activity is linear with enzyme concentration.

References:

1. Gray, H. B. et al. (1975) *Nucleic Acids Res.* 2, 1459-1492.
2. Legerski, R. J. et al. (1978) *Nucleic Acids Res.* 5, 1445-1463.
3. Wei, C. -F. et al. (1983) *J. Biol. Chem.* 258, 13506-13512.
4. Sambrook, J., Fritsch, E. F. and Maniatis, T. (1989). *Molecular Cloning: A Laboratory Manual*, (2nd ed.), (pp. 5.73-5.75). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.

CERTIFICATE OF ANALYSIS



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.