

MspI Methyltransferase



1-800-632-7799
info@neb.com
www.neb.com




M0215S 015150817081

M0215S



100 units **5,000 U/ml** **Lot: 0151508**
RECOMBINANT **Store at -20°C** **Exp: 8/17**

Methylation Site:



Description: MspI Methyltransferase recognizes the same sequence as the HpaII Methyltransferase, but modifies the external cytosine residue (C⁵) in the sequence above.

Source: An *E. coli* strain that carries the cloned MspI modification gene from *Moraxella* species (ATCC 49670)

Supplied in: 50 mM NaCl, 50 mM Tris-HCl (pH 7.5), 10 mM EDTA, 5 mM 2-mercaptoethanol, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X MspI Methyltransferase Reaction Buffer, 400X S-adenosylmethionine (32 mM).

Reaction Conditions: 1X MspI Methyltransferase Reaction Buffer, supplemented with 80 µM S-adenosylmethionine (supplied). Incubate at 37°C.

1X MspI Methyltransferase Reaction Buffer:

100 mM NaCl
50 mM Tris-HCl
10 mM EDTA
5 mM 2-mercaptoethanol
pH 7.5 @ 25°C

Protection Assay Conditions: MspI Methyltransferase is incubated with 1 µg of λ DNA in 10 µl 1X MspI Methyltransferase Reaction Buffer, supplemented with 80 µM S-adenosylmethionine, for one hour at 37°C followed by 15 minutes at 65°C. The extent of protection by MspI Methyltransferase is determined by the addition of 40 µl NEBuffer 2 supplemented with 10 mM MgCl₂ and 5 units of MspI restriction endonuclease. Incubation for MspI at 37°C is followed by analysis on agarose gels.

Unit Definition: One unit is defined as the amount of enzyme required to protect 1 µg λ DNA in 1 hour at 37°C in a total reaction volume of 10 µl against cleavage by MspI restriction endonuclease.

Quality Control Assays

16-Hour Incubation: Incubation of 10 units of MspI Methyltransferase with 1 µg of HindIII-digested λ DNA in 50 µl 1X NEBuffer 2 for 16 hours at 37°C resulted in no detectable endonuclease contamination.

Exonuclease Activity: Incubation of 10 units of MspI Methyltransferase with 1 µg sonicated [³H] DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl NEBuffer 2 [50 mM NaCl, 10 mM Tris-HCl (pH 7.9 @ 25°C), 10 mM MgCl₂, 1 mM DTT] released < 0.05% of the total radioactivity.

Storage of SAM: S-adenosylmethionine (SAM) is stored at -20°C as a 32 mM solution dissolved in 0.005 M sulfuric acid and 10% ethanol. Under these conditions SAM is stable for up to 6 months. SAM is unstable at (pH 7.5), 37°C, (1) and should be replenished in reactions incubated longer than 4 hours.

Methylation can be optimized by using fresh SAM.

Reference:

- Hoffman, J.L. (1986) *Biochemistry* 25, 4444-4449.

Companion Product:

S-adenosylmethionine (SAM)
#B9003S 0.5 ml

CERTIFICATE OF ANALYSIS

MspI Methyltransferase



1-800-632-7799
info@neb.com
www.neb.com




M0215S 015150817081

M0215S



100 units **5,000 U/ml** **Lot: 0151508**
RECOMBINANT **Store at -20°C** **Exp: 8/17**

Methylation Site:



Description: MspI Methyltransferase recognizes the same sequence as the HpaII Methyltransferase, but modifies the external cytosine residue (C⁵) in the sequence above.

Source: An *E. coli* strain that carries the cloned MspI modification gene from *Moraxella* species (ATCC 49670)

Supplied in: 50 mM NaCl, 50 mM Tris-HCl (pH 7.5), 10 mM EDTA, 5 mM 2-mercaptoethanol, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X MspI Methyltransferase Reaction Buffer, 400X S-adenosylmethionine (32 mM).

Reaction Conditions: 1X MspI Methyltransferase Reaction Buffer, supplemented with 80 µM S-adenosylmethionine (supplied). Incubate at 37°C.

1X MspI Methyltransferase Reaction Buffer:

100 mM NaCl
50 mM Tris-HCl
10 mM EDTA
5 mM 2-mercaptoethanol
pH 7.5 @ 25°C

Protection Assay Conditions: MspI Methyltransferase is incubated with 1 µg of λ DNA in 10 µl 1X MspI Methyltransferase Reaction Buffer, supplemented with 80 µM S-adenosylmethionine, for one hour at 37°C followed by 15 minutes at 65°C. The extent of protection by MspI Methyltransferase is determined by the addition of 40 µl NEBuffer 2 supplemented with 10 mM MgCl₂ and 5 units of MspI restriction endonuclease. Incubation for MspI at 37°C is followed by analysis on agarose gels.

Unit Definition: One unit is defined as the amount of enzyme required to protect 1 µg λ DNA in 1 hour at 37°C in a total reaction volume of 10 µl against cleavage by MspI restriction endonuclease.

Quality Control Assays

16-Hour Incubation: Incubation of 10 units of MspI Methyltransferase with 1 µg of HindIII-digested λ DNA in 50 µl 1X NEBuffer 2 for 16 hours at 37°C resulted in no detectable endonuclease contamination.

Exonuclease Activity: Incubation of 10 units of MspI Methyltransferase with 1 µg sonicated [³H] DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl NEBuffer 2 [50 mM NaCl, 10 mM Tris-HCl (pH 7.9 @ 25°C), 10 mM MgCl₂, 1 mM DTT] released < 0.05% of the total radioactivity.

Storage of SAM: S-adenosylmethionine (SAM) is stored at -20°C as a 32 mM solution dissolved in 0.005 M sulfuric acid and 10% ethanol. Under these conditions SAM is stable for up to 6 months. SAM is unstable at (pH 7.5), 37°C, (1) and should be replenished in reactions incubated longer than 4 hours.

Methylation can be optimized by using fresh SAM.

Reference:

- Hoffman, J.L. (1986) *Biochemistry* 25, 4444-4449.

Companion Product:

S-adenosylmethionine (SAM)
#B9003S 0.5 ml

CERTIFICATE OF ANALYSIS



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.