

HhaI Methyltransferase



1-800-632-7799
info@neb.com
www.neb.com




M0217S 006160917091

M0217S



1,000 units 25,000 U/ml Lot: 0061609
RECOMBINANT Store at -20°C Exp: 9/17

Methylation Site:



Description: HhaI Methyltransferase modifies the internal cytosine residue (C⁵) in the sequence above.

HhaI Methyltransferase



1-800-632-7799
info@neb.com
www.neb.com




M0217S 006160917091

M0217S



1,000 units 25,000 U/ml Lot: 0061609
RECOMBINANT Store at -20°C Exp: 9/17

Methylation Site:



Description: HhaI Methyltransferase modifies the internal cytosine residue (C⁵) in the sequence above.

Source: An *E. coli* strain that carries the cloned HhaI modification gene from *Haemophilus haemolyticus* (ATCC 10014)

Supplied in: 150 mM NaCl, 50 mM Tris-HCl (pH 7.5), 10 mM EDTA, 5 mM 2-mercapto-ethanol, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X HhaI Methyltransferase Reaction Buffer, 400X S-adenosylmethionine (32 mM).

Reaction Conditions: 1X HhaI Methyltransferase Reaction Buffer, 80 µM S-adenosylmethionine. Incubate at 37°C.

1X HhaI Methyltransferase Reaction Buffer:

50 mM Tris-HCl
10 mM EDTA
5 mM 2-mercaptoethanol
pH 7.5 @ 25°C

Protection Assay Conditions:

HhaI Methyltransferase is incubated with 1 µg of λ DNA in 10 µl 1X HhaI Methyltransferase Reaction Buffer, supplemented with 80 µM S-adenosylmethionine, for one hour at 37°C followed by 15 minutes at 65°C.

Source: An *E. coli* strain that carries the cloned HhaI modification gene from *Haemophilus haemolyticus* (ATCC 10014)

Supplied in: 150 mM NaCl, 50 mM Tris-HCl (pH 7.5), 10 mM EDTA, 5 mM 2-mercapto-ethanol, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:

10X HhaI Methyltransferase Reaction Buffer, 400X S-adenosylmethionine (32 mM).

Reaction Conditions: 1X HhaI Methyltransferase Reaction Buffer, 80 µM S-adenosylmethionine. Incubate at 37°C.

1X HhaI Methyltransferase Reaction Buffer:

50 mM Tris-HCl
10 mM EDTA
5 mM 2-mercaptoethanol
pH 7.5 @ 25°C

Protection Assay Conditions:

HhaI Methyltransferase is incubated with 1 µg of λ DNA in 10 µl 1X HhaI Methyltransferase Reaction Buffer, supplemented with 80 µM S-adenosylmethionine, for one hour at 37°C followed by 15 minutes at 65°C.

The extent of protection by HhaI Methyltransferase is determined by the addition of 40 µl NEBuffer 4 supplemented with 100 µg/ml BSA, 10 mM MgCl₂ and 10 units of HhaI restriction endonuclease. Incubation at 37°C for 30 minutes is followed by analysis on an agarose gel.

Unit Definition:

One unit is defined as the amount of enzyme required to protect 1 µg λ DNA in 1 hour at 37°C in a total reaction volume of 10 µl against cleavage by HhaI restriction endonuclease.

Quality Control Assays

16-Hour Incubation: Incubation of 125 units of HhaI Methyltransferase with 1 µg of HindIII-digested DNA in 50 µl 1X NEBuffer 2 for 16 hours at 37°C resulted in no detectable endonuclease contamination.

Exonuclease Activity: Incubation of 250 units of HhaI Methyltransferase with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl NEBuffer 2 [50 mM NaCl, 10 mM Tris-HCl (pH 7.9 @ 25°C), 10 mM MgCl₂, 1 mM DTT] released 0.39% of the total radioactivity.

The extent of protection by HhaI Methyltransferase is determined by the addition of 40 µl NEBuffer 4 supplemented with 100 µg/ml BSA, 10 mM MgCl₂ and 10 units of HhaI restriction endonuclease. Incubation at 37°C for 30 minutes is followed by analysis on an agarose gel.

Unit Definition:

One unit is defined as the amount of enzyme required to protect 1 µg λ DNA in 1 hour at 37°C in a total reaction volume of 10 µl against cleavage by HhaI restriction endonuclease.

Quality Control Assays

16-Hour Incubation: Incubation of 125 units of HhaI Methyltransferase with 1 µg of HindIII-digested DNA in 50 µl 1X NEBuffer 2 for 16 hours at 37°C resulted in no detectable endonuclease contamination.

Exonuclease Activity: Incubation of 250 units of HhaI Methyltransferase with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl NEBuffer 2 [50 mM NaCl, 10 mM Tris-HCl (pH 7.9 @ 25°C), 10 mM MgCl₂, 1 mM DTT] released 0.39% of the total radioactivity.

Storage of SAM: S-adenosylmethionine or SAM is stored at -20°C as a 32 mM solution dissolved in 0.005 M sulfuric acid and 10% ethanol. Under these conditions SAM is stable for up to 6 months. SAM is unstable at (pH 7.5), 37°C, (1) and should be replenished in reactions incubated longer than 4 hours.

Methylation can be optimized by using fresh SAM.

Reference:

- Hoffman, J. L. (1986) *Biochemistry* 25, 4444-4449.

Companion Product:

S-adenosylmethionine (SAM)
#B9003S 0.5 ml



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

Storage of SAM: S-adenosylmethionine or SAM is stored at -20°C as a 32 mM solution dissolved in 0.005 M sulfuric acid and 10% ethanol. Under these conditions SAM is stable for up to 6 months. SAM is unstable at (pH 7.5), 37°C, (1) and should be replenished in reactions incubated longer than 4 hours.

Methylation can be optimized by using fresh SAM.

Reference:

- Hoffman, J. L. (1986) *Biochemistry* 25, 4444-4449.

Companion Product:

S-adenosylmethionine (SAM)
#B9003S 0.5 ml



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS