

dam
Methyltransferase




M0222S 015150917091

1-800-632-7799
info@neb.com
www.neb.com

M0222S **RRR SAM**

500 units 8,000 U/ml Lot: 0151509
RECOMBINANT Store at -20°C Exp: 9/17

Methylation Site:



Description: *dam* Methyltransferase modifies the adenine residue (N⁶) in the sequence above.

dam
Methyltransferase




M0222S 015150917091

1-800-632-7799
info@neb.com
www.neb.com

M0222S **RRR SAM**

500 units 8,000 U/ml Lot: 0151509
RECOMBINANT Store at -20°C Exp: 9/17

Methylation Site:



Description: *dam* Methyltransferase modifies the adenine residue (N⁶) in the sequence above.

Source: An *E. coli* strain that carries plasmid pTP166 carrying the *dam* modification gene of *E. coli* (M. Marinus).

Supplied in: 50 mM KCl, 50 mM Tris-HCl (pH 7.5), 10 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:
10X *dam* Methyltransferase Reaction Buffer, 400X S-adenosylmethionine (32 mM).

Reaction Conditions: 1X *dam* Methyltransferase Reaction Buffer, 80 µM S-adenosylmethionine. Incubate at 37°C.

1X *dam* Methyltransferase Reaction Buffer:
50 mM Tris-HCl
10 mM EDTA
5 mM 2-mercaptoethanol
pH 7.5 @ 25°C

Source: An *E. coli* strain that carries plasmid pTP166 carrying the *dam* modification gene of *E. coli* (M. Marinus).

Supplied in: 50 mM KCl, 50 mM Tris-HCl (pH 7.5), 10 mM EDTA, 1 mM dithiothreitol, 200 µg/ml BSA and 50% glycerol.

Reagents Supplied with Enzyme:
10X *dam* Methyltransferase Reaction Buffer, 400X S-adenosylmethionine (32 mM).

Reaction Conditions: 1X *dam* Methyltransferase Reaction Buffer, 80 µM S-adenosylmethionine. Incubate at 37°C.

1X *dam* Methyltransferase Reaction Buffer:
50 mM Tris-HCl
10 mM EDTA
5 mM 2-mercaptoethanol
pH 7.5 @ 25°C

Protection Assay Conditions: *dam* Methyltransferase is incubated with 1 µg of λ DNA in 10 µl of 1X *dam* Methyltransferase Reaction Buffer, supplemented with 80 µM S-adenosylmethionine, for 1 hour at 37°C followed by 15 minutes at 65°C. The extent of protection is determined by addition of 40 µl 1X NEBuffer 3 supplemented with 10 mM MgCl₂ and 10 units of MboI restriction endonuclease. Incubation at 37°C for 1 hour is followed by analysis on an agarose gel.

Unit Definition: One unit is defined as the amount of enzyme required to protect 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 10 µl against cleavage by MboI restriction endonuclease.

Protection Assay Conditions: *dam* Methyltransferase is incubated with 1 µg of λ DNA in 10 µl of 1X *dam* Methyltransferase Reaction Buffer, supplemented with 80 µM S-adenosylmethionine, for 1 hour at 37°C followed by 15 minutes at 65°C. The extent of protection is determined by addition of 40 µl 1X NEBuffer 3 supplemented with 10 mM MgCl₂ and 10 units of MboI restriction endonuclease. Incubation at 37°C for 1 hour is followed by analysis on an agarose gel.

Unit Definition: One unit is defined as the amount of enzyme required to protect 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 10 µl against cleavage by MboI restriction endonuclease.

Quality Control Assays

16-Hour Incubation: Incubation of 60 units of *dam* Methyltransferase with 1 µg of HindIII-digested λ DNA in 50 µl 1X NEBuffer 2 for 16 hours at 37°C resulted in no detectable contamination.

Exonuclease Activity: Incubation of 120 units of *dam* Methyltransferase with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl NEBuffer 2 [50 mM NaCl, 10 mM Tris-HCl (pH 7.9 @ 25°C), 10 mM MgCl₂, 1 mM DTT] released < 0.1% of the total radioactivity.

Storage of SAM: S-adenosylmethionine (Sigma Catalog #A7007) is stored at -20°C as a 32 mM solution dissolved in 0.005 M sulfuric acid and 10% ethanol. Under these conditions SAM is stable for up to 6 months. SAM is unstable at (pH 7.5), 37°C, (1) and should be replenished in reactions incubated longer than 4 hours.

Methylation can be optimized by using fresh SAM.

Reference:

- Hoffman, J. L. (1986) *Biochemistry* 25, 4444-4449.

CERTIFICATE OF ANALYSIS

Quality Control Assays

16-Hour Incubation: Incubation of 60 units of *dam* Methyltransferase with 1 µg of HindIII-digested λ DNA in 50 µl 1X NEBuffer 2 for 16 hours at 37°C resulted in no detectable contamination.

Exonuclease Activity: Incubation of 120 units of *dam* Methyltransferase with 1 µg sonicated ³H DNA (10⁵ cpm/µg) for 4 hours at 37°C in 50 µl NEBuffer 2 [50 mM NaCl, 10 mM Tris-HCl (pH 7.9 @ 25°C), 10 mM MgCl₂, 1 mM DTT] released < 0.1% of the total radioactivity.

Storage of SAM: S-adenosylmethionine (Sigma Catalog #A7007) is stored at -20°C as a 32 mM solution dissolved in 0.005 M sulfuric acid and 10% ethanol. Under these conditions SAM is stable for up to 6 months. SAM is unstable at (pH 7.5), 37°C, (1) and should be replenished in reactions incubated longer than 4 hours.

Methylation can be optimized by using fresh SAM.

Reference:

- Hoffman, J. L. (1986) *Biochemistry* 25, 4444-4449.

CERTIFICATE OF ANALYSIS



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.