

## 9°N™ DNA Ligase



1-800-632-7799  
info@neb.com  
www.neb.com



M0238S 003150517051

# M0238S



**2,500 units**    **40,000 U/ml**    **Lot: 0031505**  
**RECOMBINANT**    **Store at -20°C**    **Exp: 5/17**

**Description:** 9°N DNA Ligase catalyzes the formation of a phosphodiester bond between juxtaposed 5' phosphate and 3' hydroxyl termini of two adjacent oligonucleotides which are hybridized to a complementary target DNA. 9°N DNA Ligase is active at elevated temperatures (45°C–90°C).

**Source:** Purified from an *E. coli* strain containing the cloned ligase gene from the extremely thermophilic marine archaea *Thermococcus* sp.(strain 9°N). The archaea was isolated from a

submarine thermal vent, at a depth of 2,500 meters, 9° north of the equator at the East Pacific Rise (1).

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 1 mM DTT, 10 mM ammonium sulfate, 200 µg/ml BSA and 50% glycerol.

### Applications:

- Allele-specific gene detection using Ligase Detection Reaction and Ligase Chain Reaction (2,4).
- Mutagenesis by incorporation of a phosphorylated oligonucleotide during PCR amplification (5).

### Reagents Supplied with Enzyme:

10X 9°N DNA Ligase Reaction Buffer:

**Reaction Conditions:** Incubate DNA and enzyme in 1X 9°N DNA Ligase Reaction Buffer at 45°C for 15 minutes or in a thermocycler with a program suited to the reaction described by Barany (1991) Genetic Disease Detection and DNA Amplification Using Cloned Thermostable Ligase. *Proc. Natl. Acad. Sci. USA* 88, 189–193. The reaction is stopped with a mixture of 50% glycerol, 50 mM EDTA, bromphenol blue.

submarine thermal vent, at a depth of 2,500 meters, 9° north of the equator at the East Pacific Rise (1).

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 1 mM DTT, 10 mM ammonium sulfate, 200 µg/ml BSA and 50% glycerol.

### Applications:

- Allele-specific gene detection using Ligase Detection Reaction and Ligase Chain Reaction (2,4).
- Mutagenesis by incorporation of a phosphorylated oligonucleotide during PCR amplification (5).

### Reagents Supplied with Enzyme:

10X 9°N DNA Ligase Reaction Buffer:

**Reaction Conditions:** Incubate DNA and enzyme in 1X 9°N DNA Ligase Reaction Buffer at 45°C for 15 minutes or in a thermocycler with a program suited to the reaction described by Barany (1991) Genetic Disease Detection and DNA Amplification Using Cloned Thermostable Ligase. *Proc. Natl. Acad. Sci. USA* 88, 189–193. The reaction is stopped with a mixture of 50% glycerol, 50 mM EDTA, bromphenol blue.

### 1X 9°N DNA Ligase Reaction Buffer:

10 mM Tris-HCl  
600 µM ATP  
2.5 mM dithiothreitol  
2.5 mM MgCl<sub>2</sub>  
0.1% Triton X-100  
(pH 7.5 @ 25°C)

### Unit Definition: (Cohesive End Unit)

One unit is defined as the amount of enzyme required to give 50% ligation of the 12-base pair cohesive ends of 1 µg of BstEII-digested λ DNA in a total reaction volume of 50 µl in 15 minutes at 45°C.

A cohesive end unit is equivalent to the nick-closing unit (1).

### Unit Assay Conditions:

1X 9°N DNA Ligase Reaction Buffer and 20 µg/ml BstEII-digested λ DNA in a 50 µl reaction. After incubation at 45°C for 15 minutes, the reaction is terminated by addition of stop dye (50% glycerol, 50 mM EDTA and bromophenol blue), heated at 70°C for 10 minutes and then loaded on a 0.7% agarose gel. Due to the presence of ligase, the cos ends of BstEII-digested λ DNA will stay together after 70°C heat treatment.

**Heat Inactivation:** No

### 1X 9°N DNA Ligase Reaction Buffer:

10 mM Tris-HCl  
600 µM ATP  
2.5 mM dithiothreitol  
2.5 mM MgCl<sub>2</sub>  
0.1% Triton X-100  
(pH 7.5 @ 25°C)

### Unit Definition: (Cohesive End Unit)

One unit is defined as the amount of enzyme required to give 50% ligation of the 12-base pair cohesive ends of 1 µg of BstEII-digested λ DNA in a total reaction volume of 50 µl in 15 minutes at 45°C.

A cohesive end unit is equivalent to the nick-closing unit (1).

### Unit Assay Conditions:

1X 9°N DNA Ligase Reaction Buffer and 20 µg/ml BstEII-digested λ DNA in a 50 µl reaction. After incubation at 45°C for 15 minutes, the reaction is terminated by addition of stop dye (50% glycerol, 50 mM EDTA and bromophenol blue), heated at 70°C for 10 minutes and then loaded on a 0.7% agarose gel. Due to the presence of ligase, the cos ends of BstEII-digested λ DNA will stay together after 70°C heat treatment.

**Heat Inactivation:** No

### Quality Control Assays

**Exonuclease Activity:** Incubation of 1,200 units for 4 hours at 37°C in 50 µl of thermostable ligase buffer containing 1 µg sonicated <sup>3</sup>H (10<sup>5</sup> cpm/µg) gave < 0.1% acid soluble counts.

**Nuclease Activity:** Incubation of HindIII fragments of λ DNA with 80 units of enzyme for 16 hours at 37°C in the recommended assay buffer without NAD does not alter the λ HindIII banding pattern on agarose gels.

**Endonuclease Activity:** Incubation of 1,500 units of enzyme for 4 hours at 37°C in 50 µl of assay buffer without NAD and 1 µg φX174 RF I DNA gave < 10% conversion to RF II

**Note: 9°N DNA Ligase is not a substitute for T4 DNA Ligase. The cohesive end unit is equivalent to the nick-closing unit of Barany et al.**

### References:

1. *Thermococcus* sp. (strain 9°N-7) isolated by Dr. Holger Jannasch, Woods Hole Oceanographic Institute, 1991.

(see other side)

CERTIFICATE OF ANALYSIS

## 9°N™ DNA Ligase



1-800-632-7799  
info@neb.com  
www.neb.com



M0238S 003150517051

# M0238S



**2,500 units**    **40,000 U/ml**    **Lot: 0031505**  
**RECOMBINANT**    **Store at -20°C**    **Exp: 5/17**

**Description:** 9°N DNA Ligase catalyzes the formation of a phosphodiester bond between juxtaposed 5' phosphate and 3' hydroxyl termini of two adjacent oligonucleotides which are hybridized to a complementary target DNA. 9°N DNA Ligase is active at elevated temperatures (45°C–90°C).

**Source:** Purified from an *E. coli* strain containing the cloned ligase gene from the extremely thermophilic marine archaea *Thermococcus* sp.(strain 9°N). The archaea was isolated from a

### Quality Control Assays

**Exonuclease Activity:** Incubation of 1,200 units for 4 hours at 37°C in 50 µl of thermostable ligase buffer containing 1 µg sonicated <sup>3</sup>H (10<sup>5</sup> cpm/µg) gave < 0.1% acid soluble counts.

**Nuclease Activity:** Incubation of HindIII fragments of λ DNA with 80 units of enzyme for 16 hours at 37°C in the recommended assay buffer without NAD does not alter the λ HindIII banding pattern on agarose gels.

**Endonuclease Activity:** Incubation of 1,500 units of enzyme for 4 hours at 37°C in 50 µl of assay buffer without NAD and 1 µg φX174 RF I DNA gave < 10% conversion to RF II

**Note: 9°N DNA Ligase is not a substitute for T4 DNA Ligase. The cohesive end unit is equivalent to the nick-closing unit of Barany et al.**

### References:

1. *Thermococcus* sp. (strain 9°N-7) isolated by Dr. Holger Jannasch, Woods Hole Oceanographic Institute, 1991.

(see other side)

CERTIFICATE OF ANALYSIS

2. Barany, F. (1991) *Proc. Natl. Acad. Sci. USA* 88, 189-193.
3. Takahashi, M. et al. (1984) *J. Biol. Chem.* 259, 10041–10047.
4. Barany, F. (1991) *The Ligase Chain Reaction in a PCR World* (pp. 5–16). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
5. Michael, S.F. (1994) *Biotechniques* 16, 411–412.



**Research Use Only: No License to Use in Any Patented Process Granted or Implied Notice to Purchaser about Limited License**

This product is designed to ligate DNA fragments at temperatures requiring a thermoactive and thermostable enzyme. The seller is aware that the product may be used in the Ligase Chain Reaction (LCR™) process covered by one or more claims of a pending patent application or issued patent assigned to Cornell Research Foundation, Inc. or Cornell Research Foundation, Inc. and the California Institute of Technology. The seller has a limited license under such patent rights, and purchase of this product includes a fully paid-up limited, non-exclusive sublicense under such intellectual property rights to use this product to perform the Ligase Chain Reaction™ process only for research and development activities and for industrial quality assurance testing. No other license has been granted expressly, impliedly or by estoppel to seller, and seller grants no other license or sublicense expressly, impliedly or by estoppel.

U.S. Publication No. 2009-0142811

Page 2 (M0238)

2. Barany, F. (1991) *Proc. Natl. Acad. Sci. USA* 88, 189-193.
3. Takahashi, M. et al. (1984) *J. Biol. Chem.* 259, 10041–10047.
4. Barany, F. (1991) *The Ligase Chain Reaction in a PCR World* (pp. 5–16). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.
5. Michael, S.F. (1994) *Biotechniques* 16, 411–412.



**Research Use Only: No License to Use in Any Patented Process Granted or Implied Notice to Purchaser about Limited License**

This product is designed to ligate DNA fragments at temperatures requiring a thermoactive and thermostable enzyme. The seller is aware that the product may be used in the Ligase Chain Reaction (LCR™) process covered by one or more claims of a pending patent application or issued patent assigned to Cornell Research Foundation, Inc. or Cornell Research Foundation, Inc. and the California Institute of Technology. The seller has a limited license under such patent rights, and purchase of this product includes a fully paid-up limited, non-exclusive sublicense under such intellectual property rights to use this product to perform the Ligase Chain Reaction™ process only for research and development activities and for industrial quality assurance testing. No other license has been granted expressly, impliedly or by estoppel to seller, and seller grants no other license or sublicense expressly, impliedly or by estoppel.

U.S. Publication No. 2009-0142811

Page 2 (M0238)