

hOGG1



1-800-632-7799
info@neb.com
www.neb.com



M0241S 003151216121

M0241S



80 units **1,600 U/ml** **Lot: 0031512**

RECOMBINANT **Store at -20°C** **Exp: 12/16**

Description: hOGG1 (α isoform) is an 8-oxoguanine DNA glycosylase which acts both as a *N*-glycosylase and an AP-lyase. The *N*-glycosylase activity releases damaged purines from double stranded DNA, generating an apurinic (AP) site. The AP-lyase activity cleaves 3' to the AP site leaving a 5' phosphate and a 3'-phospho- α , β -unsaturated aldehyde.

Some of the damaged bases recognized and removed by hOGG1 include 7, 8-dihydro-8-oxo-guanine (8-oxoguanine) when base paired with

cytosine, 8-oxoadenine when base paired with cytosine, foramidopyrimidine (fapy)-guanine and methy-fapy-guanine (1,2).

Source: An *E. coli* strain that carries the cloned human *ogg1* gene (3).

Applications:

- Single cell gel electrophoresis (Comet assay) (4,5,6)
- Alkaline elution (7)
- Alkaline unwinding (8)

Supplied in: 20 mM Tris-HCl (pH 8.0), 1 mM EDTA, 50 mM NaCl, 200 μ g/ml BSA and 50% glycerol.

Reagents supplied with Enzyme:
10X NEBuffer 2, 100X BSA.

Reaction Conditions: 1X NEBuffer 2, supplemented with 100 μ g/ml BSA. Incubate at 37°C.

1X NEBuffer 2:

50 mM NaCl
10 mM Tris-HCl
10 mM MgCl₂
1 mM DTT
pH 7.9 @ 25°C

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Unit Definition: One unit is defined as the amount of enzyme required to cleave 1 pmol of a 34-mer oligonucleotide duplex containing a single 8-oxoguanine base paired with a cytosine in 10 μ l of 1X NEBuffer 2 containing 10 pmol of substrate, supplemented with 100 μ g/ml BSA in 1 hour at 37°C.

Unit Assay Conditions: 1X NEBuffer 2 containing 10 pmol of fluorescently labeled oligonucleotide duplex, supplemented with 100 μ g/ml BSA.

Recommended dilution for the Comet Assay: 1:10² to 1:10³ (4,5,6,9). A detailed protocol can be found at www.neb.com.

Quality Control Assays

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection. BSA is added to the enzyme for stability.

16-Hour Incubation: A 50 μ l reaction containing 1 μ g of λ DNA (HindIII digest) and 16 units of hOGG1 incubated for 16 hours at 37°C resulted in DNA patterns free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Exonuclease Activity: Incubation of a 50 μ l reaction containing 8 units of hOGG1 with 1 μ g of a mixture of single and double-stranded [³H] *E. coli* DNA (10⁵ cpm/ μ g) for 4 hours at 37°C released < 0.1% of the total radioactivity.

Endonuclease Activity: Incubation of 8 units of enzyme with 1 μ g ϕ X174 RF I DNA for 4 hours at 37°C in 50 μ l reaction buffer resulted in < 5% conversion to RF II.

(See other side)

CERTIFICATE OF ANALYSIS

hOGG1



1-800-632-7799
info@neb.com
www.neb.com



M0241S 003151216121

M0241S



80 units **1,600 U/ml** **Lot: 0031512**

RECOMBINANT **Store at -20°C** **Exp: 12/16**

Description: hOGG1 (α isoform) is an 8-oxoguanine DNA glycosylase which acts both as a *N*-glycosylase and an AP-lyase. The *N*-glycosylase activity releases damaged purines from double stranded DNA, generating an apurinic (AP) site. The AP-lyase activity cleaves 3' to the AP site leaving a 5' phosphate and a 3'-phospho- α , β -unsaturated aldehyde.

Some of the damaged bases recognized and removed by hOGG1 include 7, 8-dihydro-8-oxo-guanine (8-oxoguanine) when base paired with

cytosine, 8-oxoadenine when base paired with cytosine, foramidopyrimidine (fapy)-guanine and methy-fapy-guanine (1,2).

Source: An *E. coli* strain that carries the cloned human *ogg1* gene (3).

Applications:

- Single cell gel electrophoresis (Comet assay) (4,5,6)
- Alkaline elution (7)
- Alkaline unwinding (8)

Supplied in: 20 mM Tris-HCl (pH 8.0), 1 mM EDTA, 50 mM NaCl, 200 μ g/ml BSA and 50% glycerol.

Reagents supplied with Enzyme:
10X NEBuffer 2, 100X BSA.

Reaction Conditions: 1X NEBuffer 2, supplemented with 100 μ g/ml BSA. Incubate at 37°C.

1X NEBuffer 2:

50 mM NaCl
10 mM Tris-HCl
10 mM MgCl₂
1 mM DTT
pH 7.9 @ 25°C

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Unit Definition: One unit is defined as the amount of enzyme required to cleave 1 pmol of a 34-mer oligonucleotide duplex containing a single 8-oxoguanine base paired with a cytosine in 10 μ l of 1X NEBuffer 2 containing 10 pmol of substrate, supplemented with 100 μ g/ml BSA in 1 hour at 37°C.

Unit Assay Conditions: 1X NEBuffer 2 containing 10 pmol of fluorescently labeled oligonucleotide duplex, supplemented with 100 μ g/ml BSA.

Recommended dilution for the Comet Assay: 1:10² to 1:10³ (4,5,6,9). A detailed protocol can be found at www.neb.com.

Quality Control Assays

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection. BSA is added to the enzyme for stability.

16-Hour Incubation: A 50 μ l reaction containing 1 μ g of λ DNA (HindIII digest) and 16 units of hOGG1 incubated for 16 hours at 37°C resulted in DNA patterns free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Exonuclease Activity: Incubation of a 50 μ l reaction containing 8 units of hOGG1 with 1 μ g of a mixture of single and double-stranded [³H] *E. coli* DNA (10⁵ cpm/ μ g) for 4 hours at 37°C released < 0.1% of the total radioactivity.

Endonuclease Activity: Incubation of 8 units of enzyme with 1 μ g ϕ X174 RF I DNA for 4 hours at 37°C in 50 μ l reaction buffer resulted in < 5% conversion to RF II.

(See other side)

CERTIFICATE OF ANALYSIS

Heat Inactivation: 160 units of enzyme were inactivated by incubation at 65°C for 15 minutes.

References:

1. Bjoras, M. et al. (1997). Opposite base-dependent reactions of a human base excision repair enzyme on DNA containing 7, 8-dihydro-8-oxoguanine and abasic sites. *EMBO J.* 16, 6314–6322.
2. Boiteux, S. and Radicella, J. (1999). Base excision repair of 8-hydroxyguanine protects DNA from endogenous oxidative stress. *Biochimie* 81, 59–67.
3. Radicella, J., Dherin, C. Desmze, C., Fox, M. and Boiteux, S. (1997). Cloning and characterization of *hOGG1*, a human homolog of the *OGG1* gene of *Saccharomyces cerevisiae*. *PNAS USA* 94, 8018–8015.
4. Singh, N., McCoy, M., Tice, R. and Schneider, L. (1988). A simple technique for quantitation of low levels of DNA damage in individual cells. *Experimental Cell Research* 175, 184–191.

Page 2 (M0241)

Heat Inactivation: 160 units of enzyme were inactivated by incubation at 65°C for 15 minutes.

References:

1. Bjoras, M. et al. (1997). Opposite base-dependent reactions of a human base excision repair enzyme on DNA containing 7, 8-dihydro-8-oxoguanine and abasic sites. *EMBO J.* 16, 6314–6322.
2. Boiteux, S. and Radicella, J. (1999). Base excision repair of 8-hydroxyguanine protects DNA from endogenous oxidative stress. *Biochimie* 81, 59–67.
3. Radicella, J., Dherin, C. Desmze, C., Fox, M. and Boiteux, S. (1997). Cloning and characterization of *hOGG1*, a human homolog of the *OGG1* gene of *Saccharomyces cerevisiae*. *PNAS USA* 94, 8018–8015.
4. Singh, N., McCoy, M., Tice, R. and Schneider, L. (1988). A simple technique for quantitation of low levels of DNA damage in individual cells. *Experimental Cell Research* 175, 184–191.

Page 2 (M0241)

5. Collins, A., Duthie, S. and Dobson, V. (1993). Direct enzymatic detection of endogenous oxidative base damage in human lymphocyte DNA. *Carcinogenesis* 14, 1733–1735.
6. Collins, A., Dusinska, M., Gedik, C. and Stetina, R. (1996). Oxidative damage to DNA: do we have a reliable biomarker? *Environmental Health Perspectives* 104, 465–469.
7. Pflaum, M., Will, O., Mahler, H-C. and Epe, B. (1998). DNA oxidation products determined with repair endonucleases in mammalian cells: types, basal levels and influence of cell proliferation. *Free Rad. Res.* 29, 585–594
8. Hartwig, A., Dally, H. and Schlepegrell, R. (1996). Sensitive analysis of oxidative DNA damage in mammalian cells: use of the bacterial Fpg protein in combination with alkaline unwinding. *Toxicology Letters* 88, 85–90.
9. Guthrie, E. New England Biolabs, Inc. unpublished observations



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

5. Collins, A., Duthie, S. and Dobson, V. (1993). Direct enzymatic detection of endogenous oxidative base damage in human lymphocyte DNA. *Carcinogenesis* 14, 1733–1735.
6. Collins, A., Dusinska, M., Gedik, C. and Stetina, R. (1996). Oxidative damage to DNA: do we have a reliable biomarker? *Environmental Health Perspectives* 104, 465–469.
7. Pflaum, M., Will, O., Mahler, H-C. and Epe, B. (1998). DNA oxidation products determined with repair endonucleases in mammalian cells: types, basal levels and influence of cell proliferation. *Free Rad. Res.* 29, 585–594
8. Hartwig, A., Dally, H. and Schlepegrell, R. (1996). Sensitive analysis of oxidative DNA damage in mammalian cells: use of the bacterial Fpg protein in combination with alkaline unwinding. *Toxicology Letters* 88, 85–90.
9. Guthrie, E. New England Biolabs, Inc. unpublished observations



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.