

Taq DNA Polymerase with Standard Taq Buffer



1-800-632-7799
info@neb.com
www.neb.com



M0273S 014150317031

M0273S

400 units **5,000 U/ml** **Lot: 0141503**
RECOMBINANT **Store at -20°C** **Exp: 3/17**

Description: *Taq* DNA Polymerase is a thermostable DNA polymerase that possesses a 5' → 3' polymerase activity (1,2,3) and a 5' flap endonuclease activity (4,5).

It is supplied with 10X Standard *Taq* Reaction Buffer, which is detergent-free and designed to be compatible with existing assay systems.

Source: An *E. coli* strain that carries the *Taq* DNA Polymerase gene from *Thermus aquaticus* YT-1

Application:

- PCR
- Primer Extension
- DHPLC
- Microarray Analysis
- Colony PCR

Supplied in: 100 mM KCl, 10 mM Tris-HCl (pH 7.4), 0.1 mM EDTA, 1 mM dithiothreitol, 0.5% Tween® 20, 0.5% IGEPAL® CA-630 and 50% glycerol.

Reagents Supplied with Enzyme:
10X Standard *Taq* Reaction Buffer

Reaction Conditions: 1X Standard *Taq* Reaction Buffer, DNA template, primers, 200 μM dNTPs (not included) and 1.25 units of *Taq* DNA Polymerase in a total reaction volume of 50 μl.

1X Standard *Taq* Reaction Buffer:

10 mM Tris-HCl
50 mM KCl
1.5 mM MgCl₂
pH 8.3 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 15 nmol of dNTP into acid insoluble material in 30 minutes at 75°C.

Unit Assay Conditions: 1X ThermoPol® Reaction Buffer, 200 μM dNTPs including [³H]-dTTP and 15 nM primed M13 DNA.

Heat Inactivation: No

Quality Control Assays

5 kb Lambda PCR: 25 cycles of PCR amplification of 5 ng Lambda DNA with 1.25 units of *Taq* DNA Polymerase in the presence of 200 μM dNTPs and 0.2 μM primers in Standard *Taq* Reaction Buffer results in the expected 5 kb product.

3' → 5' Exonuclease Activity: Incubation of a 20 μl reaction in ThermoPol Reaction Buffer containing a minimum of 20 units of *Taq* DNA Polymerase with 10 nM fluorescent internally labeled oligonucleotide for 30 minutes at either 37°C or 75°C yields no detectable 3' → 5' degradation as determined by capillary electrophoresis.

Endonuclease Activity: Incubation of a 50 μl reaction in ThermoPol Reaction Buffer containing a minimum of 20 units of *Taq* DNA Polymerase with 1 μg of supercoiled φX174 DNA for 4 hours at 75°C results in < 10% conversion to the nicked form as determined by agarose gel electrophoresis.

PCR

The Polymerase Chain Reaction (PCR) is a powerful and sensitive technique for DNA amplification (6). *Taq* DNA Polymerase is an enzyme widely used in PCR (7). The following guidelines are provided to ensure successful PCR using New England Biolabs' *Taq* DNA Polymerase. These guidelines cover routine PCR reactions. Amplification of templates with high GC content, high secondary structure, low template concentrations, or amplicons greater than 5 kb may require further optimization.

Reaction setup:

We recommend assembling all reaction components on ice and quickly transferring the reactions to a thermocycler preheated to the denaturation temperature (95°C).

COMPONENT	25 μl REACTION	50 μl REACTION	FINAL CONCENTRATION
10X Standard <i>Taq</i> Reaction Buffer	2.5 μl	5 μl	1X
10 mM dNTPs	0.5 μl	1 μl	200 μM
10 μM Forward Primer	0.5 μl	1 μl	0.2 μM (0.05–1 μM)
10 μM Reverse Primer	0.5 μl	1 μl	0.2 μM (0.05–1 μM)
<i>Taq</i> DNA Polymerase	0.125 μl	0.25 μl	1.25 units/50 μl PCR
Template DNA	variable	variable	<1,000 ng
Nuclease-Free Water	to 25 μl	to 50 μl	

Notes: Gently mix the reaction. Collect all liquid to the bottom of the tube by a quick spin if necessary. Overlay the sample with mineral oil if using a PCR machine without a heated lid.

Transfer PCR tubes from ice to a PCR machine with the block preheated to 95°C and begin thermocycling:

Thermocycling Conditions for a Routine PCR:

STEP	TEMP	TIME
Initial Denaturation	95°C	30 seconds
30 Cycles	95°C	15–30 seconds
	45–68°C	15–60 seconds
	68°C	1 minute/kb
Final Extension	68°C	5 minutes
Hold	4–10°C	

General Guidelines:

1. **Template:**
Use of high quality, purified DNA templates greatly enhances the success of PCR reactions. Recommended amounts of DNA template for a 50 μl reaction are as follows:

DNA	AMOUNT
Genomic	1 ng–1 μg
Plasmid or Viral	1 pg–1 ng

2. **Primers:**
Oligonucleotide primers are generally 20–40 nucleotides in length and ideally have a GC content of 40–60%. Computer programs such as Primer3 (<http://frodo.wi.mit.edu/primer3>) can be used to design or analyze primers. The final concentration of each primer in a PCR reaction may be 0.05–1 μM, typically 0.1–0.5 μM.

3. **Mg²⁺ and additives:**
Mg²⁺ concentration of 1.5–2.0 mM is optimal for most PCR products generated with *Taq* DNA Polymerase. The final Mg²⁺ concentration in 1X Standard *Taq* Reaction Buffer is 1.5 mM. This supports satisfactory amplification of most amplicons. However, Mg²⁺ can be further optimized in 0.5 or 1.0 mM increments using MgCl₂.

Amplification of some difficult targets, like GC-rich sequences, may be improved with additives, such as DMSO (8) or formamide (9).

4. **Deoxynucleotides:**
The final concentration of dNTPs is typically 200 μM of each deoxynucleotide.
5. ***Taq* DNA Polymerase Concentration:**
We generally recommend using *Taq* DNA Polymerase at a concentration of 25 units/ml (1.25 units/50 μl reaction). However, the optimal concentration of *Taq* DNA Polymerase may range from 5–50 units/ml (0.25–2.5 units/50 μl reaction) in specialized applications.
6. **Denaturation:**
An initial denaturation of 30 seconds at 95°C is sufficient for most amplicons from pure DNA templates. For difficult templates such as GC-rich sequences, a longer denaturation of 2–4 minutes at 95°C is recommended prior to PCR cycling to fully denature the template. With colony PCR, an initial 5 minute denaturation at 95°C is recommended.

During thermocycling a 15–30 second denaturation at 95°C is recommended.

7. **Annealing:**
The annealing step is typically 15–60 seconds. Annealing temperature is based on the T_m of the primer pair and is typically 45–68°C. Annealing temperatures can be optimized by doing a temperature gradient PCR starting 5°C below the calculated T_m. We recommend using NEB's T_m Calculator, available at www.neb.com/TmCalculator to determine appropriate annealing temperatures for PCR.

When primers with annealing temperatures above 60°C are used, a 2-step PCR protocol is possible (see #10).

(see other side)

8. Extension:
The recommended extension temperature is 68°C. Extension times are generally 1 minute per kb. A final extension of 5 minutes at 68°C is recommended.
9. Cycle number:
Generally, 25–35 cycles yields sufficient product. Up to 45 cycles may be required to detect low-copy-number targets.
10. 2-step PCR:
When primers with annealing temperatures above 60°C are used, a 2-step thermocycling protocol is possible.

Thermocycling Conditions for a Routine 2-Step PCR:

STEP	TEMP	TIME
Initial Denaturation	95°C	30 seconds
30 Cycles	95°C	15–30 seconds
	60–68°C	1 minute/kb
Final Extension	60–68°C	5 minutes
Hold	4–10°C	

11. PCR product:
The PCR products generated using *Taq* DNA Polymerase contain dA overhangs at the 3'-end; therefore the PCR products can be ligated to dT/dU-overhang vectors.

References:

- Chien, A., Edgar, D.B. and Trela, J.M. (1976) *J. Bact.*, 127, 1550–1557.
- Kaledin, A.S., Sliusarenko, A.G. and Gorodetskii, S.I. (1980) *Biokhimiya*, 45, 644–651.
- Lawyer, F.C. et al. (1993) *PCR Methods and Appl.*, 2, 275–287.
- Longley, M.J., Bennett, S.E. and Mosbaugh D.W. (1990) *Nucleic Acids Res.*, 18, 7317–7322.
- Lyamichev, V., Brow, M.A. and Dahlberg, J.E. (1993) *Science*, 260, 778–783.
- Saiki R.K. et al. (1985) *Science*, 230, 1350–1354.
- Powell, L.M. et al. (1987) *Cell*, 50, 831–840.
- Sun, Y., Hegamyer, G. and Colburn, N. (1993) *Biotechniques*, 15, 372–374.
- Sarkar, G., Kapelner, S. and Sommer, S.S. (1990) *Nucleic Acids Res.*, 18, 7465.

Companion Products Sold Separately:

- Standard *Taq* Reaction Buffer Pack
#B9014S 6.0 ml
- Standard *Taq* (Mg-Free) Reaction Buffer Pack
#B9015S 6.0 ml
- Magnesium Chloride (MgCl₂) Solution
#B9021S 6.0 ml
- Taq* PCR Kit
#E5000S 200 Reactions
- Taq* 2X Master Mix
#M0270S 100 Reactions
#M0270L 500 Reactions
- Quick-Load® *Taq* 2X Master Mix
#M0271S 100 Reactions
#M0271L 500 Reactions
- Taq* 5X Master Mix
#M0285S 100 Reactions
#M0285L 500 Reactions
- Deoxynucleotide Solution Set
#N0446S 25 µmol each
- Deoxynucleotide Solution Mix
#N0447S 8 µmol each
#N0447L 40 µmol each

 Annealing temperature



NEW ENGLAND BIOLABS®, QUICK-LOAD® and THERMOPOL® are registered trademarks of New England Biolabs, Inc.

IGEPAL® is a registered trademark of Rhodia Operations.

TWEEN® is a registered trademark of Uniqema Americas LLC.