

Bst DNA Polymerase, Large Fragment



1-800-632-7799
info@neb.com
www.neb.com



M0275S 051120314031

M0275S



1,600 units 8,000 U/ml Lot: 0511203
RECOMBINANT Store at -20°C Exp: 3/14

Description: *Bst* DNA Polymerase, Large Fragment is the portion of the *Bacillus stearothermophilus* DNA Polymerase protein that contains the 5' → 3' polymerase activity, but lacks 5' → 3' exonuclease activity.

Source: *Bst* DNA Polymerase, Large Fragment is prepared from an *E. coli* strain containing a genetic fusion of the *Bacillus stearothermophilus* DNA Polymerase gene, lacking the 5' → 3' exonuclease domain, and the gene coding for *E. coli* maltose binding protein (MBP). The fusion protein is purified to near homogeneity and the

MBP portion of the fusion is cleaved off *in vitro*. The remaining polymerase is purified free of MBP (1).

Applications:

- DNA sequencing through high GC regions (2,3)
- Rapid Sequencing from nanogram amounts of DNA template (4)

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 1 mM dithiothreitol, 0.1% Triton X-100 and 50% glycerol.

Reagents Supplied with Enzyme:
10X ThermoPol Reaction Buffer.

Reaction Conditions:
1X ThermoPol Reaction Buffer.
Incubate at 65°C.

1X ThermoPol Reaction Buffer:
20 mM Tris-HCl
10 mM (NH₄)₂SO₄
10 mM KCl
2 mM MgSO₄
0.1% Triton X-100
pH 8.8 @ 25°C

MBP portion of the fusion is cleaved off *in vitro*. The remaining polymerase is purified free of MBP (1).

Applications:

- DNA sequencing through high GC regions (2,3)
- Rapid Sequencing from nanogram amounts of DNA template (4)

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 1 mM dithiothreitol, 0.1% Triton X-100 and 50% glycerol.

Reagents Supplied with Enzyme:
10X ThermoPol Reaction Buffer.

Reaction Conditions:
1X ThermoPol Reaction Buffer.
Incubate at 65°C.

1X ThermoPol Reaction Buffer:
20 mM Tris-HCl
10 mM (NH₄)₂SO₄
10 mM KCl
2 mM MgSO₄
0.1% Triton X-100
pH 8.8 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 65°C.

Unit Assay Conditions: 50 mM KCl, 20 mM Tris-HCl (pH 8.8), 10 mM MgCl₂, 30 nM M13mp18 SS DNA, 70 nM M13 sequencing primer (-47) 24 mer, 200 μM dATP, 200 μM dCTP, 200 μM dGTP, 100 μM dTTP including [³H]-dTTP and 100 μg/ml BSA.

Heat Inactivation: 80°C for 20 minutes.

Quality Control Assays

16-Hour Incubation: Incubation of a 50 μl reaction in ThermoPol Reaction Buffer containing a minimum of 500 units of *Bst* DNA Polymerase, Large Fragment with 1 μg of λ DNA for 16 hours at 65°C results in no detectable change in DNA banding pattern as determined by agarose gel electrophoresis.

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 65°C.

Unit Assay Conditions: 50 mM KCl, 20 mM Tris-HCl (pH 8.8), 10 mM MgCl₂, 30 nM M13mp18 SS DNA, 70 nM M13 sequencing primer (-47) 24 mer, 200 μM dATP, 200 μM dCTP, 200 μM dGTP, 100 μM dTTP including [³H]-dTTP and 100 μg/ml BSA.

Heat Inactivation: 80°C for 20 minutes.

Quality Control Assays

16-Hour Incubation: Incubation of a 50 μl reaction in ThermoPol Reaction Buffer containing a minimum of 500 units of *Bst* DNA Polymerase, Large Fragment with 1 μg of λ DNA for 16 hours at 65°C results in no detectable change in DNA banding pattern as determined by agarose gel electrophoresis.

Exonuclease Activity: Incubation of a 50 μl reaction in ThermoPol Reaction Buffer containing a minimum of 500 units of *Bst* DNA Polymerase, Large Fragment with 1 μg of a mixture of single and double-stranded [³H] *E. coli* DNA for 4 hours at 65°C releases < 0.1% of the total radioactivity.

Endonuclease Activity: Incubation of a 50 μl reaction in ThermoPol Reaction Buffer containing a minimum of 500 units of *Bst* DNA Polymerase, Large Fragment with 1 μg of supercoiled φX174 DNA for 4 hours at 65°C results in < 10% conversion to the nicked form as determined by agarose gel electrophoresis.

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

(See other side)

CERTIFICATE OF ANALYSIS

Bst DNA Polymerase, Large Fragment



1-800-632-7799
info@neb.com
www.neb.com



M0275S 051120314031

M0275S



1,600 units 8,000 U/ml Lot: 0511203
RECOMBINANT Store at -20°C Exp: 3/14

Description: *Bst* DNA Polymerase, Large Fragment is the portion of the *Bacillus stearothermophilus* DNA Polymerase protein that contains the 5' → 3' polymerase activity, but lacks 5' → 3' exonuclease activity.

Source: *Bst* DNA Polymerase, Large Fragment is prepared from an *E. coli* strain containing a genetic fusion of the *Bacillus stearothermophilus* DNA Polymerase gene, lacking the 5' → 3' exonuclease domain, and the gene coding for *E. coli* maltose binding protein (MBP). The fusion protein is purified to near homogeneity and the

Exonuclease Activity: Incubation of a 50 μl reaction in ThermoPol Reaction Buffer containing a minimum of 500 units of *Bst* DNA Polymerase, Large Fragment with 1 μg of a mixture of single and double-stranded [³H] *E. coli* DNA for 4 hours at 65°C releases < 0.1% of the total radioactivity.

Endonuclease Activity: Incubation of a 50 μl reaction in ThermoPol Reaction Buffer containing a minimum of 500 units of *Bst* DNA Polymerase, Large Fragment with 1 μg of supercoiled φX174 DNA for 4 hours at 65°C results in < 10% conversion to the nicked form as determined by agarose gel electrophoresis.

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

(See other side)

CERTIFICATE OF ANALYSIS

Enzyme Properties

Activity in NEBuffers

ThermoPol Buffer	125%
Unit Assay Conditions	100%
NEBuffer 1	50%
NEBuffer 2	100%
NEBuffer 3	50%
NEBuffer 4	100%

NEBuffers 1, 2, 3 and 4 must be supplemented with 0.1% Triton X-100 or 100 µg/ml BSA.

Approximately 10% activity is observed in these buffers in the absence of BSA or Triton X-100.

Notes On Use: *Bst* DNA Polymerase does not exhibit 3' → 5' exonuclease activity.

100 µg/ml BSA or 0.1% Triton X-100 is required for long term storage.

Reaction temperatures above 70°C are not recommended.

Bst DNA Polymerase cannot be used for thermal cycle sequencing or PCR.

Page 2 (M0275)

Companion Products Sold Separately:

Magnesium Sulfate (MgSO ₄) Solution #B1003S	6.0 ml
ThermoPol Reaction Buffer Pack #B9004S	6.0 ml
ThermoPol II (Mg-free) Reaction Buffer Pack #B9005S	6.0 ml
ThermoPol DF (Detergent-free) Reaction Buffer Pack #B9013S	6.0 ml
Deoxynucleotide Solution Set #N0446S	25 µmol each
Deoxynucleotide Solution Mix #N0447S	8 µmol each
#N0447L	40 µmol each

References:

1. Kong, H., Aliotta, J. and Pelletier, J.J., New England Biolabs, unpublished results.
2. Griffin, H. and Griffin, A. (1994). *PCR Technology* (pp.228–229). Florida: CRC Press.
3. McClary, J. et al. (1991) *J. DNA Sequencing and Mapping* 1, 173–180.
4. Mead, D.A. et al. (1991) *BioTechniques* 11, 76–87.

U.S. Patent No. 5,814,506

Enzyme Properties

Activity in NEBuffers

ThermoPol Buffer	125%
Unit Assay Conditions	100%
NEBuffer 1	50%
NEBuffer 2	100%
NEBuffer 3	50%
NEBuffer 4	100%

NEBuffers 1, 2, 3 and 4 must be supplemented with 0.1% Triton X-100 or 100 µg/ml BSA.

Approximately 10% activity is observed in these buffers in the absence of BSA or Triton X-100.

Notes On Use: *Bst* DNA Polymerase does not exhibit 3' → 5' exonuclease activity.

100 µg/ml BSA or 0.1% Triton X-100 is required for long term storage.

Reaction temperatures above 70°C are not recommended.

Bst DNA Polymerase cannot be used for thermal cycle sequencing or PCR.

Page 2 (M0275)

Companion Products Sold Separately:

Magnesium Sulfate (MgSO ₄) Solution #B1003S	6.0 ml
ThermoPol Reaction Buffer Pack #B9004S	6.0 ml
ThermoPol II (Mg-free) Reaction Buffer Pack #B9005S	6.0 ml
ThermoPol DF (Detergent-free) Reaction Buffer Pack #B9013S	6.0 ml
Deoxynucleotide Solution Set #N0446S	25 µmol each
Deoxynucleotide Solution Mix #N0447S	8 µmol each
#N0447L	40 µmol each

References:

1. Kong, H., Aliotta, J. and Pelletier, J.J., New England Biolabs, unpublished results.
2. Griffin, H. and Griffin, A. (1994). *PCR Technology* (pp.228–229). Florida: CRC Press.
3. McClary, J. et al. (1991) *J. DNA Sequencing and Mapping* 1, 173–180.
4. Mead, D.A. et al. (1991) *BioTechniques* 11, 76–87.

U.S. Patent No. 5,814,506