

## Afu Uracil-DNA Glycosylase (UDG)



1-800-632-7799  
info@neb.com  
www.neb.com



M0279S 001120514051

# M0279S



200 units 2,000 U/ml Lot: 0011205

RECOMBINANT Store at -20°C Exp: 5/14

**Description:** A thermostable homolog of the *E. coli* Uracil-DNA Glycosylase (UDG) (1,2) from *Archaeoglobus fulgidus*. *Afu* UDG catalyzes the release of free uracil from uracil-containing DNA. *Afu* UDG efficiently hydrolyzes uracil from single-stranded or double-stranded DNA.

**Source:** Cloned and overexpressed in *E. coli*.

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.4 @ 25°C), 0.1 mM EDTA, 1 mM DTT, 0.1 mg/ml BSA and 50% glycerol.

New Reaction Buffer

## Afu Uracil-DNA Glycosylase (UDG)



1-800-632-7799  
info@neb.com  
www.neb.com



M0279S 001120514051

# M0279S



200 units 2,000 U/ml Lot: 0011205

RECOMBINANT Store at -20°C Exp: 5/14

**Description:** A thermostable homolog of the *E. coli* Uracil-DNA Glycosylase (UDG) (1,2) from *Archaeoglobus fulgidus*. *Afu* UDG catalyzes the release of free uracil from uracil-containing DNA. *Afu* UDG efficiently hydrolyzes uracil from single-stranded or double-stranded DNA.

**Source:** Cloned and overexpressed in *E. coli*.

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.4 @ 25°C), 0.1 mM EDTA, 1 mM DTT, 0.1 mg/ml BSA and 50% glycerol.

New Reaction Buffer

**Reagents Supplied with Enzyme:**  
10X ThermoPol II (Mg-free) Reaction Buffer.

**Reaction Conditions:** 1X ThermoPol II (Mg-free) Reaction Buffer. Incubate at 65°C.

**1X ThermoPol II (Mg-free) Reaction Buffer:**  
20 mM Tris-HCl  
10 mM KCl  
10 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>  
0.1% Triton X-100  
(pH 8.8 @ 25°C).

**Unit Definition:** One unit is defined as the amount of enzyme that catalyzes the release of 60 pmol of uracil per minute from double-stranded, uracil-containing DNA. Activity is measured by release of [<sup>3</sup>H]-uracil in a 50 µl reaction containing 0.2 µg DNA (10<sup>4</sup>-10<sup>5</sup> cpm/µg) in 30 minutes at 65°C.

**Unit Assay Conditions:** 1X ThermoPol II (Mg-free) Reaction Buffer, 1 unit of *Afu* Uracil-DNA Glycosylase, 0.2 µg <sup>3</sup>H-uracil DNA (10<sup>4</sup>-10<sup>5</sup> cpm/µg) for 30 minutes at 65°C in a total reaction volume of 50 µl.

**Reagents Supplied with Enzyme:**  
10X ThermoPol II (Mg-free) Reaction Buffer.

**Reaction Conditions:** 1X ThermoPol II (Mg-free) Reaction Buffer. Incubate at 65°C.

**1X ThermoPol II (Mg-free) Reaction Buffer:**  
20 mM Tris-HCl  
10 mM KCl  
10 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>  
0.1% Triton X-100  
(pH 8.8 @ 25°C).

**Unit Definition:** One unit is defined as the amount of enzyme that catalyzes the release of 60 pmol of uracil per minute from double-stranded, uracil-containing DNA. Activity is measured by release of [<sup>3</sup>H]-uracil in a 50 µl reaction containing 0.2 µg DNA (10<sup>4</sup>-10<sup>5</sup> cpm/µg) in 30 minutes at 65°C.

**Unit Assay Conditions:** 1X ThermoPol II (Mg-free) Reaction Buffer, 1 unit of *Afu* Uracil-DNA Glycosylase, 0.2 µg <sup>3</sup>H-uracil DNA (10<sup>4</sup>-10<sup>5</sup> cpm/µg) for 30 minutes at 65°C in a total reaction volume of 50 µl.

**Molecular Weight:** 22,717 daltons.

**Specific Activity:** 96 units/µg

### Quality Control Assays

**16-Hour Incubation:** A 50 µl reaction containing 1 µg of λ phage DNA in ThermoPol Reaction Buffer and 50 units of *Afu* UDG showed no degradation following overnight incubation at 65°C.

**Exonuclease Activity:** Incubation of 50 units for 4 hours at 65°C in 50 µl of ThermoPol Reaction Buffer with 1 µg <sup>3</sup>H DNA (10<sup>5</sup> cpm/µg) released < 0.1% radioactivity.

**Endonuclease Activity:** Incubation of 5 units of *Afu* UDG with 1 µg of φX174 RF I DNA for 4 hours at 65°C in 50 µl ThermoPol Reaction Buffer resulted in < 20% conversion to RF II.

**Heat Inactivation:** No

**Note:** *Afu* UDG retains 50% activity in the presence of 150 mM NaCl. *Afu* UDG retains less than 1% activity after boiling for 30 minutes in standard reaction conditions.

**Molecular Weight:** 22,717 daltons.

**Specific Activity:** 96 units/µg

### Quality Control Assays

**16-Hour Incubation:** A 50 µl reaction containing 1 µg of λ phage DNA in ThermoPol Reaction Buffer and 50 units of *Afu* UDG showed no degradation following overnight incubation at 65°C.

**Exonuclease Activity:** Incubation of 50 units for 4 hours at 65°C in 50 µl of ThermoPol Reaction Buffer with 1 µg <sup>3</sup>H DNA (10<sup>5</sup> cpm/µg) released < 0.1% radioactivity.

**Endonuclease Activity:** Incubation of 5 units of *Afu* UDG with 1 µg of φX174 RF I DNA for 4 hours at 65°C in 50 µl ThermoPol Reaction Buffer resulted in < 20% conversion to RF II.

**Heat Inactivation:** No

**Note:** *Afu* UDG retains 50% activity in the presence of 150 mM NaCl. *Afu* UDG retains less than 1% activity after boiling for 30 minutes in standard reaction conditions.

Under standard reaction conditions, uracil glycosylase inhibitor (UGI) does not inhibit *Afu* UDG.

### References:

1. Sandigursky, M. and Franklin, W. A. (1999) *Current Biology* 9, 531-534.
2. Sandigursky, M. and Franklin, W. A. (2000) *J. Biol. Chem.* 275, 19146-19149.

CERTIFICATE OF ANALYSIS

Under standard reaction conditions, uracil glycosylase inhibitor (UGI) does not inhibit *Afu* UDG.

### References:

1. Sandigursky, M. and Franklin, W. A. (1999) *Current Biology* 9, 531-534.
2. Sandigursky, M. and Franklin, W. A. (2000) *J. Biol. Chem.* 275, 19146-19149.

CERTIFICATE OF ANALYSIS