

## Exonuclease I (*E. coli*)



1-800-632-7799  
info@neb.com  
www.neb.com



M0293S 022151117111

# M0293S



3,000 units 20,000 U/ml Lot: 0221511

RECOMBINANT Store at -20°C Exp: 11/17

**Description:** Catalyzes the removal of nucleotides from single-stranded DNA in the 3' to 5' direction.

**Source:** An *E. coli* strain that carries the cloned *Exo I* gene from *E. coli* NM554

### Applications:

- Exonuclease I degrades excess single-stranded primer oligonucleotide from a reaction mixture containing double-stranded extension products.

## Exonuclease I (*E. coli*)



1-800-632-7799  
info@neb.com  
www.neb.com



M0293S 022151117111

# M0293S



3,000 units 20,000 U/ml Lot: 0221511

RECOMBINANT Store at -20°C Exp: 11/17

**Description:** Catalyzes the removal of nucleotides from single-stranded DNA in the 3' to 5' direction.

**Source:** An *E. coli* strain that carries the cloned *Exo I* gene from *E. coli* NM554

### Applications:

- Exonuclease I degrades excess single-stranded primer oligonucleotide from a reaction mixture containing double-stranded extension products.

**Storage Conditions:** 100 mM NaCl, 10 mM Tris-HCl (pH 7.5), 0.5 mM EDTA, 5 mM 2-mercaptoethanol, 100 µg/ml BSA and 50% glycerol.

**Reagents Supplied with Enzyme:**  
10X Exonuclease I Reaction Buffer

**Reaction Conditions:** 1X Exonuclease I Reaction Buffer. Incubate at 37°C.

**1X Exonuclease I Reaction Buffer:**  
67 mM Glycine-KOH  
6.7 mM MgCl<sub>2</sub>  
10 mM 2-mercaptoethanol  
(pH 9.5 @ 25°C)

**Unit Definition:** One unit is defined as the amount of enzyme that will catalyze the release of 10 nmol of acid-soluble nucleotide in 30 minutes at 37°C under standard reaction conditions.

**Unit Assay Conditions:** 67 mM Glycine-KOH (pH 9.5), 6.7 mM MgCl<sub>2</sub>, 10 mM 2-mercaptoethanol, and 0.17 mg/ml single-stranded [<sup>3</sup>H]-DNA.

**Heat Inactivation:** 80°C for 20 minutes.

**Storage Conditions:** 100 mM NaCl, 10 mM Tris-HCl (pH 7.5), 0.5 mM EDTA, 5 mM 2-mercaptoethanol, 100 µg/ml BSA and 50% glycerol.

**Reagents Supplied with Enzyme:**  
10X Exonuclease I Reaction Buffer

**Reaction Conditions:** 1X Exonuclease I Reaction Buffer. Incubate at 37°C.

**1X Exonuclease I Reaction Buffer:**  
67 mM Glycine-KOH  
6.7 mM MgCl<sub>2</sub>  
10 mM 2-mercaptoethanol  
(pH 9.5 @ 25°C)

**Unit Definition:** One unit is defined as the amount of enzyme that will catalyze the release of 10 nmol of acid-soluble nucleotide in 30 minutes at 37°C under standard reaction conditions.

**Unit Assay Conditions:** 67 mM Glycine-KOH (pH 9.5), 6.7 mM MgCl<sub>2</sub>, 10 mM 2-mercaptoethanol, and 0.17 mg/ml single-stranded [<sup>3</sup>H]-DNA.

**Heat Inactivation:** 80°C for 20 minutes.

### Quality Control Assays

**Double-stranded Endonuclease:** Incubation of 100 units of Exonuclease I with 1 µg φX174 RF I DNA for 16 hours at 37°C in 50 µl reaction buffer resulted in < 5% conversion to RF II.

**Single-stranded Endonuclease:** Incubation of 100 units of Exonuclease I with 1 µg of M13mp18 single-stranded DNA for 16 hours at 37°C in a 50 µl reaction resulted in no decrease in the amount of closed circular DNA as determined by agarose gel electrophoresis.

**Double-stranded Exonuclease:** Incubation of 50 units of enzyme with 0.2 µg <sup>3</sup>H DNA (9.0 x 10<sup>3</sup> cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.5% radioactivity.

**Physical Purity:** Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection. BSA is added to the enzyme for stability.

### Quality Control Assays

**Double-stranded Endonuclease:** Incubation of 100 units of Exonuclease I with 1 µg φX174 RF I DNA for 16 hours at 37°C in 50 µl reaction buffer resulted in < 5% conversion to RF II.

**Single-stranded Endonuclease:** Incubation of 100 units of Exonuclease I with 1 µg of M13mp18 single-stranded DNA for 16 hours at 37°C in a 50 µl reaction resulted in no decrease in the amount of closed circular DNA as determined by agarose gel electrophoresis.

**Double-stranded Exonuclease:** Incubation of 50 units of enzyme with 0.2 µg <sup>3</sup>H DNA (9.0 x 10<sup>3</sup> cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.5% radioactivity.

**Physical Purity:** Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection. BSA is added to the enzyme for stability.

### References:

- Lehman and Nussbaum (1964) *J. Biol. Chem.* 239, 2628.
- Kusher et al. (1971) *Proc. Natl. Acad. Sci. USA* 68, 824.
- Kusher et al. (1972) *Proc. Natl. Acad. Sci. USA* 69, 1366.
- Goldmark and Linn (1972) *J. Biol. Chem.* 247, 184.
- Rosamond et al. (1979) *J. Biol. Chem.* 254, 8646.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

### References:

- Lehman and Nussbaum (1964) *J. Biol. Chem.* 239, 2628.
- Kusher et al. (1971) *Proc. Natl. Acad. Sci. USA* 68, 824.
- Kusher et al. (1972) *Proc. Natl. Acad. Sci. USA* 69, 1366.
- Goldmark and Linn (1972) *J. Biol. Chem.* 247, 184.
- Rosamond et al. (1979) *J. Biol. Chem.* 254, 8646.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS