

Tth Endonuclease IV



1-800-632-7799
info@neb.com
www.neb.com



M0294S 001151117111

M0294S



500 units **10,000 U/ml** **Lot: 0011511**
RECOMBINANT **Store at -20°** **Exp: 11/17**

Description: *Tth* Endonuclease IV is a thermostable apurinic/aprimidinic (AP) endonuclease from *Thermus thermophilus*. *Tth* Endo IV will hydrolyze an AP site at the first phosphodiester bond 5' to the lesion leaving a 3' hydroxyl and a deoxyribose 5'-phosphate at the 5' terminus. The enzyme also has a 3'-diesterase activity.

Source: An *E. coli* strain that carries the cloned *Thermus thermophilus* endonuclease IV gene.

More Units

Applications:

- Alkaline elution (1)
- Alkaline unwinding (2)

Supplied in: 100 mM KCl, 10 mM Tris-HCl (pH 7.4 @ 25°C), 0.1 mM EDTA, 1 mM DTT, 0.1% Triton X-100 and 50% glycerol.

Reagents Supplied with Enzyme:
10X ThermoPol Reaction Buffer.

Reaction Conditions: 1X ThermoPol Reaction Buffer. Incubate at 65°C.

1X ThermoPol Reaction Buffer:

10 mM KCl
20 mM Tris-HCl
10 mM (NH₄)₂SO₄
2 mM MgSO₄
0.1% Triton X-100
pH 8.8 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to cleave 1 pmol of a 60-mer oligonucleotide duplex containing a single AP site* in a total reaction volume of 10 µl in 1 hour at 65°C.

Applications:

- Alkaline elution (1)
- Alkaline unwinding (2)

Supplied in: 100 mM KCl, 10 mM Tris-HCl (pH 7.4 @ 25°C), 0.1 mM EDTA, 1 mM DTT, 0.1% Triton X-100 and 50% glycerol.

Reagents Supplied with Enzyme:
10X ThermoPol Reaction Buffer.

Reaction Conditions: 1X ThermoPol Reaction Buffer. Incubate at 65°C.

1X ThermoPol Reaction Buffer:

10 mM KCl
20 mM Tris-HCl
10 mM (NH₄)₂SO₄
2 mM MgSO₄
0.1% Triton X-100
pH 8.8 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme required to cleave 1 pmol of a 60-mer oligonucleotide duplex containing a single AP site* in a total reaction volume of 10 µl in 1 hour at 65°C.

* An AP site is created by treating 10 pmol of a 60-mer oligonucleotide duplex containing a single uracil residue with 1 unit of Uracil-DNA Glycosylase (UDG) for 2 minutes at 37°C.

Diluent Compatibility: Diluent D
100 mM KCl, 10 mM Tris-HCl (pH 7.4 @ 25°C), 0.1 mM EDTA, 1 mM DTT, 0.1% Triton X-100 and 50 % glycerol.

Unit Assay Conditions: 1X ThermoPol Reaction Buffer containing 5 pmol of fluorescently labeled oligonucleotide duplex in a total reaction volume of 10 µl.

Quality Control Assays

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

16-Hour Incubation: A 50 µl reaction containing 1 µg of λ DNA (HindIII digest) and 100 units of *Tth* Endonuclease IV incubated for 16 hours at 37°C resulted in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

* An AP site is created by treating 10 pmol of a 60-mer oligonucleotide duplex containing a single uracil residue with 1 unit of Uracil-DNA Glycosylase (UDG) for 2 minutes at 37°C.

Diluent Compatibility: Diluent D
100 mM KCl, 10 mM Tris-HCl (pH 7.4 @ 25°C), 0.1 mM EDTA, 1 mM DTT, 0.1% Triton X-100 and 50 % glycerol.

Unit Assay Conditions: 1X ThermoPol Reaction Buffer containing 5 pmol of fluorescently labeled oligonucleotide duplex in a total reaction volume of 10 µl.

Quality Control Assays

Physical Purity: Purified to > 95% homogeneity as determined by SDS-PAGE analysis using Coomassie Blue detection.

16-Hour Incubation: A 50 µl reaction containing 1 µg of λ DNA (HindIII digest) and 100 units of *Tth* Endonuclease IV incubated for 16 hours at 37°C resulted in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Exonuclease Activity: Incubation of a 50 µl reaction containing 100 units of *Tth* Endonuclease IV in NEBuffer 1 with 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA (200,000 cpm/µg) for 4 hours at 37°C released < 0.1% of the total radioactivity.

Endonuclease Activity: Incubation of a 50 µl reaction containing 100 units of *Tth* Endonuclease IV with 1 µg φX174 RF I DNA for 4 hours at 37°C resulted in < 10% conversion to RF II as determined by agarose gel electrophoresis.

Heat Inactivation: no.

Usage Note: Enzyme stability above 80°C is assured by adding ZnCl₂ to a final concentration of 25 µM in the reaction.

References:

1. Pflaum, M. et al. (1998) *Free Rad. Res.* 29, 585-594.
2. Hartwig, A. et al. (1996) *Toxicology Letters* 88, 85-90.

CERTIFICATE OF ANALYSIS

Tth Endonuclease IV



1-800-632-7799
info@neb.com
www.neb.com



M0294S 001151117111

M0294S



500 units **10,000 U/ml** **Lot: 0011511**
RECOMBINANT **Store at -20°** **Exp: 11/17**

Description: *Tth* Endonuclease IV is a thermostable apurinic/aprimidinic (AP) endonuclease from *Thermus thermophilus*. *Tth* Endo IV will hydrolyze an AP site at the first phosphodiester bond 5' to the lesion leaving a 3' hydroxyl and a deoxyribose 5'-phosphate at the 5' terminus. The enzyme also has a 3'-diesterase activity.

Source: An *E. coli* strain that carries the cloned *Thermus thermophilus* endonuclease IV gene.

More Units

Exonuclease Activity: Incubation of a 50 µl reaction containing 100 units of *Tth* Endonuclease IV in NEBuffer 1 with 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA (200,000 cpm/µg) for 4 hours at 37°C released < 0.1% of the total radioactivity.

Endonuclease Activity: Incubation of a 50 µl reaction containing 100 units of *Tth* Endonuclease IV with 1 µg φX174 RF I DNA for 4 hours at 37°C resulted in < 10% conversion to RF II as determined by agarose gel electrophoresis.

Heat Inactivation: no.

Usage Note: Enzyme stability above 80°C is assured by adding ZnCl₂ to a final concentration of 25 µM in the reaction.

References:

1. Pflaum, M. et al. (1998) *Free Rad. Res.* 29, 585-594.
2. Hartwig, A. et al. (1996) *Toxicology Letters* 88, 85-90.

CERTIFICATE OF ANALYSIS



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.