

Topoisomerase I (*E. coli*)



1-800-632-7799
info@neb.com
www.neb.com



M0301S 008150615061

M0301S



100 units **5,000 U/ml** **Lot: 0081506**

RECOMBINANT **Store at -20°C** **Exp: 6/16**

Description: Topoisomerase I (*E. coli*) catalyzes the relaxation of negatively supercoiled DNA. Topoisomerase I has also been implicated in knotting and unknotting DNA (1) and in linking complementary rings of single-stranded DNA into double-stranded rings (2). The intact holoenzyme is a 97 kDa protein.

Source: An *E. coli* strain containing the cloned topA gene from *E. coli*.

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.5), 35 mM (NH₄)₂SO₄, 0.1 mM EDTA, 1 mM DTT and 50% glycerol.

Reagents Supplied with Enzyme:
10X CutSmart™ Buffer.

Reaction Conditions: 1X CutSmart Buffer.
Incubate at 37°C.

1X CutSmart Buffer:

50 mM potassium acetate
20 mM Tris acetate
10 mM magnesium acetate
100 µg/ml BSA
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme that catalyzes the relaxation of > 95% of 0.5 µg of negatively supercoiled pUC19 RF I DNA in a total reaction volume of 25 µl in 15 minutes at 37°C. DNA supercoiling is assessed by agarose gel electrophoresis in the absence of ethidium bromide.

Quality Assurance: Purified free of contaminating exonucleases and endonucleases.

Quality Control Assays:

16-Hour Incubation: A 50 µl reaction containing 1 µg of λ DNA/HindIII fragments and 50 units of Topoisomerase I incubated for 16 hours resulted in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Exonuclease Activity: Incubation of 50 units of Topoisomerase I with 1 µg sonicated single and doubled-stranded [³H] DNA (200,000 cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.5% radioactivity.

Endonuclease Activity: Incubation of 40 units of Topoisomerase I with 1 µg pUC19 DNA for 4 hours at 37°C in 50 µl reaction buffer resulted in < 10% conversion to RF II, as determined by agarose gel electrophoresis in the presence of ethidium bromide.

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1	25%
NEBuffer 2	50%
NEBuffer 3	0%
NEBuffer 4	100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 0.5 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 40 units of Topoisomerase I were inactivated by incubation at 65°C for 20 minutes.

References:

1. Liu, L.F. et al. (1976) *J. Mol. Biol.* 106, 439–452.
2. Kirkegaard, K. and Wang, J.C. (1978) *Nucleic Acids Res.* 5, 3811–3820.

(see other side)

CERTIFICATE OF ANALYSIS

Topoisomerase I (*E. coli*)



1-800-632-7799
info@neb.com
www.neb.com



M0301S 008150615061

M0301S



100 units **5,000 U/ml** **Lot: 0081506**

RECOMBINANT **Store at -20°C** **Exp: 6/16**

Description: Topoisomerase I (*E. coli*) catalyzes the relaxation of negatively supercoiled DNA. Topoisomerase I has also been implicated in knotting and unknotting DNA (1) and in linking complementary rings of single-stranded DNA into double-stranded rings (2). The intact holoenzyme is a 97 kDa protein.

Source: An *E. coli* strain containing the cloned topA gene from *E. coli*.

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.5), 35 mM (NH₄)₂SO₄, 0.1 mM EDTA, 1 mM DTT and 50% glycerol.

Reagents Supplied with Enzyme:
10X CutSmart™ Buffer.

Reaction Conditions: 1X CutSmart Buffer.
Incubate at 37°C.

1X CutSmart Buffer:

50 mM potassium acetate
20 mM Tris acetate
10 mM magnesium acetate
100 µg/ml BSA
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme that catalyzes the relaxation of > 95% of 0.5 µg of negatively supercoiled pUC19 RF I DNA in a total reaction volume of 25 µl in 15 minutes at 37°C. DNA supercoiling is assessed by agarose gel electrophoresis in the absence of ethidium bromide.

Quality Assurance: Purified free of contaminating exonucleases and endonucleases.

Quality Control Assays:

16-Hour Incubation: A 50 µl reaction containing 1 µg of λ DNA/HindIII fragments and 50 units of Topoisomerase I incubated for 16 hours resulted in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Exonuclease Activity: Incubation of 50 units of Topoisomerase I with 1 µg sonicated single and doubled-stranded [³H] DNA (200,000 cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.5% radioactivity.

Endonuclease Activity: Incubation of 40 units of Topoisomerase I with 1 µg pUC19 DNA for 4 hours at 37°C in 50 µl reaction buffer resulted in < 10% conversion to RF II, as determined by agarose gel electrophoresis in the presence of ethidium bromide.

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1	25%
NEBuffer 2	50%
NEBuffer 3	0%
NEBuffer 4	100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 0.5 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 40 units of Topoisomerase I were inactivated by incubation at 65°C for 20 minutes.

References:

1. Liu, L.F. et al. (1976) *J. Mol. Biol.* 106, 439–452.
2. Kirkegaard, K. and Wang, J.C. (1978) *Nucleic Acids Res.* 5, 3811–3820.

(see other side)

CERTIFICATE OF ANALYSIS



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

CUTSMART™ is a trademark of New England Biolabs, Inc.

Page 2 (M0301)



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

CUTSMART™ is a trademark of New England Biolabs, Inc.

Page 2 (M0301)