

Topoisomerase I (*E. coli*)



1-800-632-7799
info@neb.com
www.neb.com



M0301S 008151216121

M0301S



100 units **5,000 U/ml** **Lot: 0081512**

RECOMBINANT **Store at -20°C** **Exp: 12/16**

Description: Topoisomerase I (*E. coli*) catalyzes the relaxation of negatively supercoiled DNA. Topoisomerase I has also been implicated in knotting and unknotting DNA (1) and in linking complementary rings of single-stranded DNA into double-stranded rings (2). The intact holoenzyme is a 97 kDa protein.

Source: An *E. coli* strain containing the cloned topA gene from *E. coli*.

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.5), 35 mM (NH₄)₂SO₄, 0.1 mM EDTA, 1 mM DTT and 50% glycerol.

Reagents Supplied with Enzyme:
10X CutSmart® Buffer.

Reaction Conditions: 1X CutSmart Buffer.
Incubate at 37°C.

1X CutSmart Buffer:

50 mM potassium acetate
20 mM Tris acetate
10 mM magnesium acetate
100 µg/ml BSA
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme that catalyzes the relaxation of > 95% of 0.5 µg of negatively supercoiled pUC19 RF I DNA in a total reaction volume of 25 µl in 15 minutes at 37°C. DNA supercoiling is assessed by agarose gel electrophoresis in the absence of ethidium bromide.

Quality Assurance: Purified free of contaminating exonucleases and endonucleases.

Quality Control Assays:

16-Hour Incubation: A 50 µl reaction containing 1 µg of λ DNA/HindIII fragments and 50 units of Topoisomerase I incubated for 16 hours resulted in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Exonuclease Activity: Incubation of 50 units of Topoisomerase I with 1 µg sonicated single and doubled-stranded [³H] DNA (200,000 cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.5% radioactivity.

Endonuclease Activity: Incubation of 40 units of Topoisomerase I with 1 µg pUC19 DNA for 4 hours at 37°C in 50 µl reaction buffer resulted in < 10% conversion to RF II, as determined by agarose gel electrophoresis in the presence of ethidium bromide.

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1	25%
NEBuffer 2	50%
NEBuffer 3	0%
NEBuffer 4	100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 0.5 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 40 units of Topoisomerase I were inactivated by incubation at 65°C for 20 minutes.

References:

1. Liu, L.F. et al. (1976) *J. Mol. Biol.* 106, 439–452.
2. Kirkegaard, K. and Wang, J.C. (1978) *Nucleic Acids Res.* 5, 3811–3820.

(see other side)

CERTIFICATE OF ANALYSIS

Topoisomerase I (*E. coli*)



1-800-632-7799
info@neb.com
www.neb.com



M0301S 008151216121

M0301S



100 units **5,000 U/ml** **Lot: 0081512**

RECOMBINANT **Store at -20°C** **Exp: 12/16**

Description: Topoisomerase I (*E. coli*) catalyzes the relaxation of negatively supercoiled DNA. Topoisomerase I has also been implicated in knotting and unknotting DNA (1) and in linking complementary rings of single-stranded DNA into double-stranded rings (2). The intact holoenzyme is a 97 kDa protein.

Source: An *E. coli* strain containing the cloned topA gene from *E. coli*.

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.5), 35 mM (NH₄)₂SO₄, 0.1 mM EDTA, 1 mM DTT and 50% glycerol.

Reagents Supplied with Enzyme:
10X CutSmart® Buffer.

Reaction Conditions: 1X CutSmart Buffer.
Incubate at 37°C.

1X CutSmart Buffer:

50 mM potassium acetate
20 mM Tris acetate
10 mM magnesium acetate
100 µg/ml BSA
pH 7.9 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme that catalyzes the relaxation of > 95% of 0.5 µg of negatively supercoiled pUC19 RF I DNA in a total reaction volume of 25 µl in 15 minutes at 37°C. DNA supercoiling is assessed by agarose gel electrophoresis in the absence of ethidium bromide.

Quality Assurance: Purified free of contaminating exonucleases and endonucleases.

Quality Control Assays:

16-Hour Incubation: A 50 µl reaction containing 1 µg of λ DNA/HindIII fragments and 50 units of Topoisomerase I incubated for 16 hours resulted in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Exonuclease Activity: Incubation of 50 units of Topoisomerase I with 1 µg sonicated single and doubled-stranded [³H] DNA (200,000 cpm/µg) for 4 hours at 37°C in 50 µl reaction buffer released < 0.5% radioactivity.

Endonuclease Activity: Incubation of 40 units of Topoisomerase I with 1 µg pUC19 DNA for 4 hours at 37°C in 50 µl reaction buffer resulted in < 10% conversion to RF II, as determined by agarose gel electrophoresis in the presence of ethidium bromide.

Enzyme Properties

Activity in NEBuffers:

NEBuffer 1	25%
NEBuffer 2	50%
NEBuffer 3	0%
NEBuffer 4	100%

When using a buffer other than the optimal (supplied) NEBuffer, it may be necessary to add more enzyme to achieve complete digestion.

Survival in a Reaction: A minimum of 0.5 unit is required to digest 1 µg of substrate DNA in 16 hours.

Heat Inactivation: 40 units of Topoisomerase I were inactivated by incubation at 65°C for 20 minutes.

References:

1. Liu, L.F. et al. (1976) *J. Mol. Biol.* 106, 439–452.
2. Kirkegaard, K. and Wang, J.C. (1978) *Nucleic Acids Res.* 5, 3811–3820.

(see other side)

CERTIFICATE OF ANALYSIS



NEW ENGLAND BIOLABS® and CUTSMART® are registered trademarks of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.



NEW ENGLAND BIOLABS® and CUTSMART® are registered trademarks of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.