

DNA Gyrase (*E. coli*)



1-800-632-7799
info@neb.com
www.neb.com



M0306S 002150916091

M0306S



100 units Lot: 0021509 Exp: 9/16
5,000 U/ml Store at **-80°C**

Description: DNA Gyrase (*E. coli*) is a Type II topoisomerase that catalyzes the introduction of negative supercoils in DNA in the presence of ATP. The gyrase holoenzyme is a heterotetramer made up of 2 gyrA (97 kDa) subunits and 2 gyrB (90 kDa) subunits.

Source: An *E. coli* strain containing the cloned and expressed gyrA and gyrB genes.

Note: Store at -80°C

DNA Gyrase (*E. coli*)



1-800-632-7799
info@neb.com
www.neb.com



M0306S 002150916091

M0306S



100 units Lot: 0021509 Exp: 9/16
5,000 U/ml Store at **-80°C**

Description: DNA Gyrase (*E. coli*) is a Type II topoisomerase that catalyzes the introduction of negative supercoils in DNA in the presence of ATP. The gyrase holoenzyme is a heterotetramer made up of 2 gyrA (97 kDa) subunits and 2 gyrB (90 kDa) subunits.

Source: An *E. coli* strain containing the cloned and expressed gyrA and gyrB genes.

Note: Store at -80°C

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 2 mM DTT and 50% glycerol.

Reagents Supplied with Enzyme:
5X DNA Gyrase (*E. coli*) Reaction Buffer

Reaction Conditions: 1X DNA Gyrase (*E. coli*) Reaction Buffer. Incubate at 37°C.

1X DNA Gyrase (*E. coli*) Reaction Buffer:

35 mM Tris-HCl
24 mM KCl
4 mM MgCl₂
2 mM DTT
1.75 mM ATP
5 mM spermidine
0.1 mg/ml BSA
6.5% glycerol
pH 7.5 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme that catalyzes the conversion of DNA Gyrase (*E. coli*) Substrate (NEB #N0471) to > 95% of 0.5 µg of supercoiled plasmid in a total reaction volume of 30 µl in 30 minutes at 37°C. DNA supercoiling is assessed by agarose gel electrophoresis in the absence of ethidium bromide.

Supplied in: 50 mM KCl, 10 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 2 mM DTT and 50% glycerol.

Reagents Supplied with Enzyme:
5X DNA Gyrase (*E. coli*) Reaction Buffer

Reaction Conditions: 1X DNA Gyrase (*E. coli*) Reaction Buffer. Incubate at 37°C.

1X DNA Gyrase (*E. coli*) Reaction Buffer:

35 mM Tris-HCl
24 mM KCl
4 mM MgCl₂
2 mM DTT
1.75 mM ATP
5 mM spermidine
0.1 mg/ml BSA
6.5% glycerol
pH 7.5 @ 25°C

Unit Definition: One unit is defined as the amount of enzyme that catalyzes the conversion of DNA Gyrase (*E. coli*) Substrate (NEB #N0471) to > 95% of 0.5 µg of supercoiled plasmid in a total reaction volume of 30 µl in 30 minutes at 37°C. DNA supercoiling is assessed by agarose gel electrophoresis in the absence of ethidium bromide.

Quality Control Assays:

16-Hour Incubation: A 50 µl reaction containing 1 µg of λDNA (HindIII digest) and 25 units of DNA Gyrase (*E. coli*) for 16 hours at 37°C resulted in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Exonuclease Activity: Incubation of a 50 µl reaction containing 25 units of DNA Gyrase (*E. coli*) with 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA (200,000 cpm/µg) for 4 hours at 37°C in released < 0.5% of the total radioactivity.

Endonuclease Activity: Incubation of 25 units of DNA Gyrase (*E. coli*) with 1 µg pUC19 DNA for 4 hours at 37°C in 50 µl reaction buffer resulted in < 10% conversion to RF II.

Enzyme Properties

Survival in a Reaction: A minimum of 0.13 unit is required to convert 0.5 µg of substrate DNA to supercoil plasmid in a total reaction volume of 30 µl in 16 hours.

Quality Control Assays:

16-Hour Incubation: A 50 µl reaction containing 1 µg of λDNA (HindIII digest) and 25 units of DNA Gyrase (*E. coli*) for 16 hours at 37°C resulted in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Exonuclease Activity: Incubation of a 50 µl reaction containing 25 units of DNA Gyrase (*E. coli*) with 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA (200,000 cpm/µg) for 4 hours at 37°C in released < 0.5% of the total radioactivity.

Endonuclease Activity: Incubation of 25 units of DNA Gyrase (*E. coli*) with 1 µg pUC19 DNA for 4 hours at 37°C in 50 µl reaction buffer resulted in < 10% conversion to RF II.

Enzyme Properties

Survival in a Reaction: A minimum of 0.13 unit is required to convert 0.5 µg of substrate DNA to supercoil plasmid in a total reaction volume of 30 µl in 16 hours.

Heat Inactivation: 25 units of DNA Gyrase (*E. coli*) were inactivated by incubation at 65°C for 20 minutes.

Note: The 5X DNA Gyrase Reaction Buffer should be stored at -80°C to maintain optimal stability of components.

References:

1. Adachi, T. et al. (1987) *Nucleic Acids Res.* 15, 771-784.
2. Higgins, N.P. et al. (1978) *PNAS* 75, 1773-1777.
3. Swanberg, S.L. and Wang, J.C. (1987) *J. Mol. Biol.* 197, 729-736.

Companion Product:

DNA Gyrase (*E. coli*) Substrate
#N0471S 20 µg

CERTIFICATE OF ANALYSIS

Heat Inactivation: 25 units of DNA Gyrase (*E. coli*) were inactivated by incubation at 65°C for 20 minutes.

Note: The 5X DNA Gyrase Reaction Buffer should be stored at -80°C to maintain optimal stability of components.

References:

1. Adachi, T. et al. (1987) *Nucleic Acids Res.* 15, 771-784.
2. Higgins, N.P. et al. (1978) *PNAS* 75, 1773-1777.
3. Swanberg, S.L. and Wang, J.C. (1987) *J. Mol. Biol.* 197, 729-736.

Companion Product:

DNA Gyrase (*E. coli*) Substrate
#N0471S 20 µg

CERTIFICATE OF ANALYSIS



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.