

5' Deadenylase



1-800-632-7799
info@neb.com
www.neb.com



M0331S 003160818081

M0331S



2,500 units 50,000 U/ml Lot: 0031608
RECOMBINANT Store at -20°C Exp: 8/18

Description: Yeast 5' Deadenylase is a member of the HIT (histidine triad) family proteins and specifically a member of the Hint branch. It is the yeast orthologue of aprataxin. Mutations in human aprataxin have been known to be involved in the neurological disorder known as ataxia oculomotor apraxia-1 (1). The human protein has been shown to resolve abortive ligation intermediates by removing the AMP at the 5' end of DNA (AMP-DNA hydrolase activity) (2). It also repairs DNA damage at 3' ends by removing

3'-phosphate and 3'-phosphoglycolate (3). Human aprataxin acts on small molecules as well, nucleotide polyphosphates such as diadenosine tetraphosphate (AppppA) as well as lysyl-AMP.

The 5' Deadenylase is encoded by *HNT3* gene of *S. cerevisiae*. NEB has shown this protein is capable of deadenylation from 5' end of DNA and RNA, leaving the phosphate at 5' end. It also cleaves AppppA into ATP and AMP. Its activity on lysyl-AMP is not detectable.

Source: Purified from an *E. coli* strain carrying a plasmid encoding 5' Deadenylase from *S. cerevisiae*

Supplied in: 100 mM NaCl, 10 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 1 mM dithiothreitol, 0.1% Triton X-100 and 50% glycerol.

Applications:

- Deadenylation of 5' end of DNA and RNA
- Aprataxin-dependent DNA repair assay
- Analysis of dinucleoside tetraphosphate

Reagents Supplied with Enzyme:

10X NEBuffer 1

3'-phosphate and 3'-phosphoglycolate (3). Human aprataxin acts on small molecules as well, nucleotide polyphosphates such as diadenosine tetraphosphate (AppppA) as well as lysyl-AMP.

The 5' Deadenylase is encoded by *HNT3* gene of *S. cerevisiae*. NEB has shown this protein is capable of deadenylation from 5' end of DNA and RNA, leaving the phosphate at 5' end. It also cleaves AppppA into ATP and AMP. Its activity on lysyl-AMP is not detectable.

Source: Purified from an *E. coli* strain carrying a plasmid encoding 5' Deadenylase from *S. cerevisiae*

Supplied in: 100 mM NaCl, 10 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 1 mM dithiothreitol, 0.1% Triton X-100 and 50% glycerol.

Applications:

- Deadenylation of 5' end of DNA and RNA
- Aprataxin-dependent DNA repair assay
- Analysis of dinucleoside tetraphosphate

Reagents Supplied with Enzyme:

10X NEBuffer 1

Reaction Conditions: 1X NEBuffer 1, 5–50 pmol adenylated DNA (AMP-DNA) in 20 μ l.

Incubate at 30°C.

1X NEBuffer 1:

10 mM Bis-Tris-Propane-HCl
10 mM MgCl₂
1 mM dithiothreitol
pH 7.0 @ 25°C

Unit Definition: One unit is defined as the amount enzyme required to remove 10 pmoles of AMP from a 5'-adenylated DNA oligo in 10 minutes at 30°C.

Protocol for a Typical Deadenylation Reaction:

Combine the following components in sterile microfuge tube:

10X NEBuffer 1	2 μ l
5–50 pmol 5'-adenylated DNA or RNA	variable
H ₂ O	variable
5' Deadenylase (50 units/ μ l)	1 μ l
Total reaction volume	20 μ l

Incubate at 30°C for 0.5–1 hour

Reaction Conditions: 1X NEBuffer 1, 5–50 pmol adenylated DNA (AMP-DNA) in 20 μ l.

Incubate at 30°C.

1X NEBuffer 1:

10 mM Bis-Tris-Propane-HCl
10 mM MgCl₂
1 mM dithiothreitol
pH 7.0 @ 25°C

Unit Definition: One unit is defined as the amount enzyme required to remove 10 pmoles of AMP from a 5'-adenylated DNA oligo in 10 minutes at 30°C.

Protocol for a Typical Deadenylation Reaction:

Combine the following components in sterile microfuge tube:

10X NEBuffer 1	2 μ l
5–50 pmol 5'-adenylated DNA or RNA	variable
H ₂ O	variable
5' Deadenylase (50 units/ μ l)	1 μ l
Total reaction volume	20 μ l

Incubate at 30°C for 0.5–1 hour

Quality Control Assays

RNase Assay: Incubation of 50 units of 5' Deadenylase with 40 ng of RNA transcript in 10 μ l for 4 hours at 37°C resulted in no detectable degradation of RNA as determined by denaturing PAGE analysis.

Endonuclease Activity: Incubation of 50 units of 5' Deadenylase with 1 μ g of supercoiled plasmid in 10 μ l for 4 hours at 37°C produced less than 10% nicked or linear molecules as determined by agarose gel electrophoresis.

Exonuclease Assay: Incubation of a 50 μ l reaction containing 100 units of 5' Deadenylase with 1 μ g of a mixture of single and double-stranded [³H] *E. coli* DNA (10⁵ cpm/ μ g) for 4 hours at 37°C released < 0.5% of the total radioactivity.

5' DNA Phosphatase Assay: Incubation of 50 units of 5' Deadenylase with 10 pmol of 5'-pDNA oligo in 10 μ l for 2 hours at 37°C resulted in no detectable dephosphorylation as determined by denaturing PAGE analysis.

(See other side)

CERTIFICATE OF ANALYSIS

5' Deadenylase



1-800-632-7799
info@neb.com
www.neb.com



M0331S 003160818081

M0331S



2,500 units 50,000 U/ml Lot: 0031608
RECOMBINANT Store at -20°C Exp: 8/18

Description: Yeast 5' Deadenylase is a member of the HIT (histidine triad) family proteins and specifically a member of the Hint branch. It is the yeast orthologue of aprataxin. Mutations in human aprataxin have been known to be involved in the neurological disorder known as ataxia oculomotor apraxia-1 (1). The human protein has been shown to resolve abortive ligation intermediates by removing the AMP at the 5' end of DNA (AMP-DNA hydrolase activity) (2). It also repairs DNA damage at 3' ends by removing

3'-phosphate and 3'-phosphoglycolate (3). Human aprataxin acts on small molecules as well, nucleotide polyphosphates such as diadenosine tetraphosphate (AppppA) as well as lysyl-AMP.

The 5' Deadenylase is encoded by *HNT3* gene of *S. cerevisiae*. NEB has shown this protein is capable of deadenylation from 5' end of DNA and RNA, leaving the phosphate at 5' end. It also cleaves AppppA into ATP and AMP. Its activity on lysyl-AMP is not detectable.

Source: Purified from an *E. coli* strain carrying a plasmid encoding 5' Deadenylase from *S. cerevisiae*

Supplied in: 100 mM NaCl, 10 mM Tris-HCl (pH 7.5), 0.1 mM EDTA, 1 mM dithiothreitol, 0.1% Triton X-100 and 50% glycerol.

Applications:

- Deadenylation of 5' end of DNA and RNA
- Aprataxin-dependent DNA repair assay
- Analysis of dinucleoside tetraphosphate

Reagents Supplied with Enzyme:

10X NEBuffer 1

Reaction Conditions: 1X NEBuffer 1, 5–50 pmol adenylated DNA (AMP-DNA) in 20 μ l.

Incubate at 30°C.

1X NEBuffer 1:

10 mM Bis-Tris-Propane-HCl
10 mM MgCl₂
1 mM dithiothreitol
pH 7.0 @ 25°C

Unit Definition: One unit is defined as the amount enzyme required to remove 10 pmoles of AMP from a 5'-adenylated DNA oligo in 10 minutes at 30°C.

Protocol for a Typical Deadenylation Reaction:

Combine the following components in sterile microfuge tube:

10X NEBuffer 1	2 μ l
5–50 pmol 5'-adenylated DNA or RNA	variable
H ₂ O	variable
5' Deadenylase (50 units/ μ l)	1 μ l
Total reaction volume	20 μ l

Incubate at 30°C for 0.5–1 hour

Quality Control Assays

RNase Assay: Incubation of 50 units of 5' Deadenylase with 40 ng of RNA transcript in 10 μ l for 4 hours at 37°C resulted in no detectable degradation of RNA as determined by denaturing PAGE analysis.

Endonuclease Activity: Incubation of 50 units of 5' Deadenylase with 1 μ g of supercoiled plasmid in 10 μ l for 4 hours at 37°C produced less than 10% nicked or linear molecules as determined by agarose gel electrophoresis.

Exonuclease Assay: Incubation of a 50 μ l reaction containing 100 units of 5' Deadenylase with 1 μ g of a mixture of single and double-stranded [³H] *E. coli* DNA (10⁵ cpm/ μ g) for 4 hours at 37°C released < 0.5% of the total radioactivity.

5' DNA Phosphatase Assay: Incubation of 50 units of 5' Deadenylase with 10 pmol of 5'-pDNA oligo in 10 μ l for 2 hours at 37°C resulted in no detectable dephosphorylation as determined by denaturing PAGE analysis.

(See other side)

CERTIFICATE OF ANALYSIS

Heat Inactivation: 70°C for 20 minutes.

Notes:

5´-adenylated DNA could be dsDNA, ssDNA and nicked DNA with the adenylate residue at the 5´ terminus of the nick. Deadenylation is more efficient with dsDNA and nicked DNA.

References:

- 1) Le Ber, I. et al. (2003) *Brain*, 126, 2761.
- 2) Ahel, I. et al. (2006) *Nature*, 443, 713.
- 3) Takahashi T. et al. (2007) *Nucl. Acids Res.*, 35, 3797.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

Heat Inactivation: 70°C for 20 minutes.

Notes:

5´-adenylated DNA could be dsDNA, ssDNA and nicked DNA with the adenylate residue at the 5´ terminus of the nick. Deadenylation is more efficient with dsDNA and nicked DNA.

References:

- 1) Le Ber, I. et al. (2003) *Brain*, 126, 2761.
- 2) Ahel, I. et al. (2006) *Nature*, 443, 713.
- 3) Takahashi T. et al. (2007) *Nucl. Acids Res.*, 35, 3797.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.