

**OneTaq® Quick-Load®  
2X Master Mix with  
GC Buffer**



1-800-632-7799  
info@neb.com  
www.neb.com



**M0487S**



**100 reactions (50 µl vol) Lot: 0061205**

**RECOMBINANT Store at -20°C Exp: 5/13**

**Description:** OneTaq Quick-Load 2X Master Mix with GC Buffer is an optimized, ready-to-use blend of Taq and Deep Vent<sub>R</sub> DNA Polymerases ideally suited to PCR applications from GC-rich templates, including pure DNA solutions, bacterial colonies and cDNA products. The 3'→5' exonuclease activity of Deep Vent DNA Polymerase increases the fidelity and robust amplification of Taq DNA Polymerase (1). The convenient quick-load master mix formulation contains dNTPs, MgSO<sub>4</sub>, buffer components and stabilizers as well as two commonly used tracking dyes for DNA gels. On a 1% agarose gel in 1X TBE, Xylene Cyanol FF migrates at ~4 kb and Tartrazine migrates at ~10 bp. Both dyes are present in concentrations that do not mask co-migrating DNA bands.

**Source:** An *E. coli* strain that carries the Taq DNA Polymerase gene from *Thermus aquaticus* YT-1 and an *E. coli* strain that carries the Deep Vent<sub>R</sub> DNA Polymerase gene from *Pyrococcus* species GB-D.

**Applications:**

- GC-rich PCR
- High Sensitivity PCR
- High Throughput PCR
- Colony PCR
- Long PCR (up to ~6 kb genomic)

**Reagents Supplied with Enzyme:**

OneTaq High GC Enhancer

**Reaction Conditions:** 1X OneTaq Quick-Load Master Mix with GC Buffer, DNA template and primers in a total reaction volume of 50 µl.

**1X OneTaq Quick-Load Master Mix with GC Buffer:**

- 80 mM Tris-SO<sub>4</sub> (pH 9.2)
- 2 mM MgSO<sub>4</sub>
- 20 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>
- 0.2 mM dNTPs
- 5% Glycerol
- 5% DMSO
- 0.06% IGEPAL® CA-630
- 0.05% Tween-20
- Xylene Cyanol FF
- Tartrazine
- 25 units/ml OneTaq DNA Polymerase

**OneTaq High GC Enhancer:**

- 10 mM Tris-HCl
- 25% DMSO
- 25% Glycerol, pH 9.2 @ 25°C

**Unit Definition:** One unit is defined as the amount of enzyme that will incorporate 15 nmol of dNTP into acid insoluble material in 30 minutes at 75°C.

**Unit Assay Conditions:** 1X ThermoPol Reaction Buffer, 200 µM dNTPs including [<sup>3</sup>H]-dTTP and 200 µg/ml activated Calf Thymus DNA.

**Heat Inactivation:** No

**Quality Control Assays**


**Buffer-dependent GC-rich (> 65% GC) PCR:**

30 cycles of PCR amplification of 10 ng of human genomic DNA with 1X OneTaq Quick-Load Master Mix with GC Buffer in a 25 µl reaction in the presence of 0.2 µM primers resulted in the buffer-dependent production of the 737 bp GC-rich product.

**Enhancer-dependent High GC (> 70% GC) PCR:**

30 cycles of PCR amplification of 10 ng of human genomic DNA with 1X OneTaq Quick-Load Master Mix with GC Buffer in a 25 µl reaction in the presence of 0.2 µM primers and 20% OneTaq High GC Enhancer resulted in the enhancer-dependent production of the 627 bp high GC product.

**Note:** Product specifications for individual components in the OneTaq Quick-Load 2X Master Mix with GC Buffer are available separately.



Amplification of a selection of sequences with varying GC content from human genomic DNA using OneTaq DNA Polymerase. GC content is indicated above gel. Marker M is the 1 kb DNA Ladder (NEB #N3232).

**PCR**

The Polymerase Chain Reaction (PCR) is a powerful and sensitive technique for DNA amplification (2). Taq DNA Polymerase is an enzyme widely used in PCR (3). The following guidelines are provided to ensure successful PCR using New England Biolabs' OneTaq Quick-Load 2X Master Mix with GC Buffer. These guidelines cover routine PCR reactions. Specialized applications may require further optimization.

**Reaction Setup:**

We recommend assembling all reaction components on ice and quickly transferring the reactions to a thermocycler preheated to the denaturation temperature (94°C).

COMPONENT	25 µl REACTION	50 µl REACTION	FINAL CONCENTRATION
OneTaq Quick-Load 2X Master Mix with GC Buffer	12.5 µl	25 µl	1X
(OneTaq High GC Enhancer, optional)*	(2.5–5 µl)	(5–10 µl)	(10–20%)
10 µM Forward Primer	0.5 µl	1 µl	0.2 µM
10 µM Reverse Primer	0.5 µl	1 µl	0.2 µM
Template DNA	variable	variable	<1,000 ng
Nuclease-Free Water	to 25 µl	to 50 µl	

\*For extremely difficult or high GC amplicons, the addition of 10–20% OneTaq High GC Enhancer may improve amplification.

**Notes:** Gently mix the reaction. Collect all liquid to the bottom of the tube by a quick spin if necessary. Overlay the sample with mineral oil if using a PCR machine without a heated lid.

Transfer PCR tubes to a PCR machine and begin thermocycling:

**Thermocycling Conditions for a Routine PCR:**

STEP	TEMP	TIME
Initial Denaturation	94°C	30 seconds
30 Cycles	94°C	15–30 seconds
	45–68°C	15–60 seconds
	68°C	1 minute/kb
Final Extension	68°C	5 minutes
Hold	4–10°C	

**General Guidelines:**

1. **Template:**  
Use of high quality, purified DNA templates greatly enhances the success of PCR reactions. Recommended amounts of DNA template for a 50 µl reaction are as follows:

DNA	AMOUNT
Genomic	1 ng–1 µg
Plasmid or Viral	1 pg–1 ng

2. **Primers:**  
Oligonucleotide primers are generally 20–40 nucleotides in length and ideally have a GC content of 40–60%. Computer programs such as Primer3 (<http://frodo.wi.mit.edu/primer3>) can be used to design or analyze primers. The final concentration of each primer in a PCR reaction may be 0.05–1 µM, typically 0.2 µM.
3. **Mg<sup>++</sup> and Additives:**  
Mg<sup>++</sup> concentration of 1.5–2.0 mM is optimal for most PCR products generated with OneTaq DNA Polymerase. The final Mg<sup>++</sup> concentration in 1X OneTaq Quick-Load Master Mix with GC Buffer is 2 mM. This supports satisfactory amplification of most amplicons. However, Mg<sup>++</sup> can be further optimized in 0.2 mM increments using MgSO<sub>4</sub> (sold separately).

Amplification of extremely difficult targets may be improved by the addition of 10–20% OneTaq High GC Enhancer (included).

(see other side)

4. Denaturation:  
An initial denaturation of 30 seconds at 94°C is sufficient to amplify most targets from pure DNA templates. For difficult templates, a longer denaturation of 2–4 minutes at 94°C is recommended prior to PCR cycling to fully denature the template. With colony PCR, an initial 2–5 minute denaturation at 94°C is recommended.

During thermocycling a 15–30 second denaturation at 94°C is recommended.

5. Annealing:  
The annealing step is typically 15–60 seconds. Annealing temperature is based on the  $T_m$  of the primer pair and is typically 45–68°C. Annealing temperatures can be optimized by doing a temperature gradient PCR starting 5°C below the calculated  $T_m$ .

6. Extension:  
The recommended extension temperature is 68°C. Extension times are generally 1 minute per kb. A final extension of 5 minutes at 68°C is recommended.

7. Cycle Number:  
Generally, 25–35 cycles yields sufficient product. Up to 45 cycles may be required to detect low copy number targets.

8. 2-step PCR:  
When primers with annealing temperatures of 68°C or above are used, a 2-step thermocycling protocol (combining annealing and extension into one step) is possible.

9. PCR Product:  
The majority of the PCR products generated using *OneTaq* DNA Polymerase contain dA overhangs at the 3' end; therefore the PCR products can be ligated to dT/dU-overhang vectors.

#### Notes:

*OneTaq* Quick-Load 2X Master Mix with GC Buffer is stable for fifteen freeze-thaw cycles when stored at –20°C.

*OneTaq* Quick-Load 2X Master Mix with GC Buffer is also stable for one month at 4°C, so for frequent use, an aliquot may be kept at 4°C.

#### References:

1. Barnes, W.M. (1994) *Proc. Natl. Acad. Sci. USA*, 91, 2216–2220.
2. Saiki R.K. et al. (1985) *Science*, 230, 1350–1354.
3. Powell, L.M. et al. (1987) *Cell*, 50, 831–840.

#### Companion Products Sold Separately:

Magnesium Sulfate ( $MgSO_4$ ) Solution  
#B1003S 6.0 ml

*OneTaq*™ Quick-Load® 2X Master Mix  
with Standard Buffer  
#M0486S 100 Reactions  
#M0486L 500 Reactions

*OneTaq*™ Hot Start Quick-Load® 2X Master Mix  
with Standard Buffer  
#M0488S 100 Reactions  
#M0488L 500 Reactions

*OneTaq*™ Hot Start Quick-Load® 2X Master Mix  
with GC Buffer  
#M0489S 100 Reactions  
#M0489L 500 Reactions

*OneTaq*™ DNA Polymerase  
#M0480S 200 units  
#M0480L 1,000 units  
#M0480X 5,000 units

*OneTaq*™ Hot Start DNA Polymerase  
#M0481S 200 units  
#M0481L 1,000 units  
#M0481X 5,000 units

#### Notice to Purchaser:

Purchase of this product provides the purchaser with a non-exclusive license for research purposes only. The purchase of this product also conveys to the purchaser only the limited, non-transferable right to use the purchased quantity of the product for the purchaser's own research by the purchaser only under the following U.S. patent claims and foreign counterpart patent claims: U.S. Patent No. 5,436,149 (claims 6-16). No rights are granted to the purchaser to sell, modify for resale or otherwise transfer this product, either alone or as a component of another product, to any third party. Takara Bio reserves all other rights, and this product may not be used in any manner other than as provided herein. For information on obtaining a license to use this product for purposes other than research, please contact Takara Bio Inc., Seta 3-4-1, Otsu, Shiga 520-2193, Japan (Fax +81-77-453-9254).

IGEPAL® is a registered trademark of Rhodia Operations.