

## Tth RecA



1-800-632-7799  
info@neb.com  
www.neb.com



M2402S 071150817081

# M2402S



**100 µg 1 mg/ml Lot: 0711508**  
**RECOMBINANT Store at -20°C Exp: 8/17**

**Description:** Tth RecA is a RecA homolog isolated from *Thermus thermophilus*. It has a ssDNA-dependent ATPase activity at an optimal temperature between 65–75°C. The extreme thermostability makes Tth RecA ideal for molecular biology applications that require an elevated temperature condition, such as nucleic acid amplification and sequencing.

**Source:** An *E. coli* strain that carries the cloned RecA gene from *Thermus thermophilus*. Developed and produced by Biohelix Corporation, a New England Biolabs-affiliated company.

## Tth RecA



1-800-632-7799  
info@neb.com  
www.neb.com



M2402S 071150817081

# M2402S



**100 µg 1 mg/ml Lot: 0711508**  
**RECOMBINANT Store at -20°C Exp: 8/17**

**Description:** Tth RecA is a RecA homolog isolated from *Thermus thermophilus*. It has a ssDNA-dependent ATPase activity at an optimal temperature between 65–75°C. The extreme thermostability makes Tth RecA ideal for molecular biology applications that require an elevated temperature condition, such as nucleic acid amplification and sequencing.

**Source:** An *E. coli* strain that carries the cloned RecA gene from *Thermus thermophilus*. Developed and produced by Biohelix Corporation, a New England Biolabs-affiliated company.

### Applications:

- Visualization of DNA structures with electron microscopy (1)
- Site-directed mutagenesis through D-loop (2,3)
- Screening of DNA libraries using RecA-probe filaments (4,5)
- Targeted cleavage of DNA (6)

Supplied in: 10 mM Tris-HCl (pH 7.5 @ 25°C), 100 mM KCl, 0.1 mM EDTA, 1 mM DTT, 0.1% Triton X-100 and 50% glycerol.

**Unit Definition:** Sold by mass of pure protein as determined by OD<sub>280</sub>.

**Molecular Weight:** 36 kDa.

**Quality Assurance:** RecA is purified free of contaminating endonucleases and exonucleases. Each lot is tested for single-stranded DNA-dependent ATPase activity and is visually determined to be > 95% pure on an SDS-polyacrylamide gel.

### Applications:

- Visualization of DNA structures with electron microscopy (1)
- Site-directed mutagenesis through D-loop (2,3)
- Screening of DNA libraries using RecA-probe filaments (4,5)
- Targeted cleavage of DNA (6)

Supplied in: 10 mM Tris-HCl (pH 7.5 @ 25°C), 100 mM KCl, 0.1 mM EDTA, 1 mM DTT, 0.1% Triton X-100 and 50% glycerol.

**Unit Definition:** Sold by mass of pure protein as determined by OD<sub>280</sub>.

**Molecular Weight:** 36 kDa.

**Quality Assurance:** RecA is purified free of contaminating endonucleases and exonucleases. Each lot is tested for single-stranded DNA-dependent ATPase activity and is visually determined to be > 95% pure on an SDS-polyacrylamide gel.

### Quality Control Assays

**Exonuclease Activity:** Incubation of 20 µg RecA for 4 hours at 65°C in 50 µl reaction buffer containing 50 mM potassium acetate (pH 7.9), 20 mM tris-acetate, 10 mM magnesium acetate and 1 mM dithiothreitol (pH 7.9 at 25°C), with a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA (200,000 cpm/µg) released < 0.1% of the total radioactivity.

**Endonuclease Activity:** Incubation of 2 µg RecA for 4 hours at 65°C in 50 µl reaction buffer containing 50 mM potassium acetate (pH 7.9), 20 mM tris-acetate, 10 mM magnesium acetate and 1 mM dithiothreitol (pH 7.9 at 25°C), with 1 µg φX174 RF I DNA gave < 20% conversion to RF II.

**Nuclease Activity:** Incubation of 20 µg RecA for 16 hours at 65°C in 50 µl of reaction buffer containing 50 mM potassium acetate (pH 7.9), 20 mM tris-acetate, 10 mM magnesium acetate and 1 mM dithiothreitol (pH 7.9 at 25°C), with 1 µg λ DNA yielded a clear and sharp band on an agarose gel.

### Quality Control Assays

**Exonuclease Activity:** Incubation of 20 µg RecA for 4 hours at 65°C in 50 µl reaction buffer containing 50 mM potassium acetate (pH 7.9), 20 mM tris-acetate, 10 mM magnesium acetate and 1 mM dithiothreitol (pH 7.9 at 25°C), with a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA (200,000 cpm/µg) released < 0.1% of the total radioactivity.

**Endonuclease Activity:** Incubation of 2 µg RecA for 4 hours at 65°C in 50 µl reaction buffer containing 50 mM potassium acetate (pH 7.9), 20 mM tris-acetate, 10 mM magnesium acetate and 1 mM dithiothreitol (pH 7.9 at 25°C), with 1 µg φX174 RF I DNA gave < 20% conversion to RF II.

**Nuclease Activity:** Incubation of 20 µg RecA for 16 hours at 65°C in 50 µl of reaction buffer containing 50 mM potassium acetate (pH 7.9), 20 mM tris-acetate, 10 mM magnesium acetate and 1 mM dithiothreitol (pH 7.9 at 25°C), with 1 µg λ DNA yielded a clear and sharp band on an agarose gel.

**Notes On Use:** Tth RecA is active in any polymerase buffer. Add 400 ng of the Tth RecA per 50 µl reaction.

### References:

1. Wasserman, S.A. and Cozzarelli, N.R. (1985) *Proc. Natl. Acad. Sci. USA*, 82, 1079–1083.
2. Biet, E. et al. (2001) *Biochemistry*, 40, 1779–1786.
3. Shortle, D. et al. (1980) *Proc. Natl. Acad. Sci. USA*, 77, 5375–5379.
4. Rigas, B. et al. (1986) *Proc. Natl. Acad. Sci. USA*, 83, 9591–9595.
5. Honigberg, S.M. et al. (1986) *Proc. Natl. Acad. Sci. USA*, 83, 9586–9590.
6. Koob, M. et al. (1992) *Nucleic Acids Res.* 20, 5831–5836.
7. Shigemori, Y. et al. (2005) *Nucleic Acids Res.* e126.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS

**Notes On Use:** Tth RecA is active in any polymerase buffer. Add 400 ng of the Tth RecA per 50 µl reaction.

### References:

1. Wasserman, S.A. and Cozzarelli, N.R. (1985) *Proc. Natl. Acad. Sci. USA*, 82, 1079–1083.
2. Biet, E. et al. (2001) *Biochemistry*, 40, 1779–1786.
3. Shortle, D. et al. (1980) *Proc. Natl. Acad. Sci. USA*, 77, 5375–5379.
4. Rigas, B. et al. (1986) *Proc. Natl. Acad. Sci. USA*, 83, 9591–9595.
5. Honigberg, S.M. et al. (1986) *Proc. Natl. Acad. Sci. USA*, 83, 9586–9590.
6. Koob, M. et al. (1992) *Nucleic Acids Res.* 20, 5831–5836.
7. Shigemori, Y. et al. (2005) *Nucleic Acids Res.* e126.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

CERTIFICATE OF ANALYSIS