

# Histone H1<sup>0</sup> Human, Recombinant



1-800-632-7799  
info@neb.com  
www.neb.com



M2501S 006150817081

## M2501S



**100 µg**      **1.0 mg/ml**      **Lot: 0061508**  
**RECOMBINANT**    **Store at -20°C**    **Exp: 8/17**


**Description:** Histone H1 acts on the linker region of polynucleosome DNA to condense the chromatin into structures of ~30 nm (1). It is not necessary for octamer or nucleosome core particle formation. Eight different Histone H1 proteins have been identified in the human genome (2). Histone H1<sup>0</sup> is a non replication-dependent histone that is highly expressed in cells that have terminally differentiated (3).

**Source:** An *E. coli* strain that carries a plasmid encoding the human histone H1 gene, H1F0 or H1FV. (Genbank accession number: X03473)

Supplied in: 20 mM Sodium Phosphate (pH 7.0), 300 mM NaCl and 1 mM EDTA.

**Note:** The protein concentration (1 mg/ml, 48 µM) is calculated using the molar extinction coefficient for Histone H1 (3840) and its absorbance at 280 nm (4,5). 1.0 A<sub>280</sub> units = 5.4 mg/ml

Synonyms: Histone H1.0, Histone H1(0), Histone H1'




**SDS-PAGE analysis of Histone H1<sup>0</sup> Human, Recombinant.**  
Lane 1 and 7: NEB Protein Ladder (NEB #P7703), Lanes 2 thru 6: 0.5, 1.0, 2.0, 5.0, 10.0 µg Histone H1<sup>0</sup> Human, Recombinant.

### Quality Control Assays:

**SDS-PAGE:** 0.5, 1.0, 2.0, 5.0, 10.0 µg of Histone H1<sup>0</sup> Human, Recombinant were loaded on a 10–20% Tris-Glycine SDS-PAGE gel and stained with Coomassie Blue. The calculated molecular weight is 20731.53 Da. Its apparent molecular weight on 10–20% Tris-Glycine SDS-PAGE gel is ~27 kDa.

**Mass Spectrometry:** The mass of purified Histone H1<sup>0</sup> Human, Recombinant is 20731.74 Da as determined by ESI-TOF MS (Electrospray



ESI-TOF Analysis of Histone H1<sup>0</sup> Human, Recombinant.

Ionization-Time of Flight Mass Spectrometry). The average mass calculated from primary sequence is 20731.53 Da. This confirms the protein identity of the histone. There is a small percentage of histone H1<sup>0</sup> with a mass of 20863.27 which is a +131 Da difference from the major species. This correlates to histone H1<sup>0</sup> with an intact N-terminal methionine (6).

**Protease Assay:** After incubation of 5 µg of Histone H1<sup>0</sup> Human, Recombinant with a standard mixture of proteases for 4 hours at 37°C, no proteolytic activity could be detected by SDS-PAGE.

**Exonuclease Assay:** Incubation of a 50 µl reaction containing 10 µg of Histone H1<sup>0</sup> Human, Recombinant with 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA (200,000 cpm/µg) for 4 hours at 37°C released < 0.1% of the total radioactivity.

(see other side)

CERTIFICATE OF ANALYSIS

# Histone H1<sup>0</sup> Human, Recombinant



1-800-632-7799  
info@neb.com  
www.neb.com



M2501S 006150817081

## M2501S



**100 µg**      **1.0 mg/ml**      **Lot: 0061508**  
**RECOMBINANT**    **Store at -20°C**    **Exp: 8/17**


**Description:** Histone H1 acts on the linker region of polynucleosome DNA to condense the chromatin into structures of ~30 nm (1). It is not necessary for octamer or nucleosome core particle formation. Eight different Histone H1 proteins have been identified in the human genome (2). Histone H1<sup>0</sup> is a non replication-dependent histone that is highly expressed in cells that have terminally differentiated (3).

**Source:** An *E. coli* strain that carries a plasmid encoding the human histone H1 gene, H1F0 or H1FV. (Genbank accession number: X03473)

Supplied in: 20 mM Sodium Phosphate (pH 7.0), 300 mM NaCl and 1 mM EDTA.

**Note:** The protein concentration (1 mg/ml, 48 µM) is calculated using the molar extinction coefficient for Histone H1 (3840) and its absorbance at 280 nm (4,5). 1.0 A<sub>280</sub> units = 5.4 mg/ml

Synonyms: Histone H1.0, Histone H1(0), Histone H1'




**SDS-PAGE analysis of Histone H1<sup>0</sup> Human, Recombinant.**  
Lane 1 and 7: NEB Protein Ladder (NEB #P7703), Lanes 2 thru 6: 0.5, 1.0, 2.0, 5.0, 10.0 µg Histone H1<sup>0</sup> Human, Recombinant.

### Quality Control Assays:

**SDS-PAGE:** 0.5, 1.0, 2.0, 5.0, 10.0 µg of Histone H1<sup>0</sup> Human, Recombinant were loaded on a 10–20% Tris-Glycine SDS-PAGE gel and stained with Coomassie Blue. The calculated molecular weight is 20731.53 Da. Its apparent molecular weight on 10–20% Tris-Glycine SDS-PAGE gel is ~27 kDa.

**Mass Spectrometry:** The mass of purified Histone H1<sup>0</sup> Human, Recombinant is 20731.74 Da as determined by ESI-TOF MS (Electrospray



ESI-TOF Analysis of Histone H1<sup>0</sup> Human, Recombinant.

Ionization-Time of Flight Mass Spectrometry). The average mass calculated from primary sequence is 20731.53 Da. This confirms the protein identity of the histone. There is a small percentage of histone H1<sup>0</sup> with a mass of 20863.27 which is a +131 Da difference from the major species. This correlates to histone H1<sup>0</sup> with an intact N-terminal methionine (6).

**Protease Assay:** After incubation of 5 µg of Histone H1<sup>0</sup> Human, Recombinant with a standard mixture of proteases for 4 hours at 37°C, no proteolytic activity could be detected by SDS-PAGE.

**Exonuclease Assay:** Incubation of a 50 µl reaction containing 10 µg of Histone H1<sup>0</sup> Human, Recombinant with 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA (200,000 cpm/µg) for 4 hours at 37°C released < 0.1% of the total radioactivity.

(see other side)

CERTIFICATE OF ANALYSIS

**Endonuclease Assay:** Incubation of a 50 µl reaction containing 10 µg of Histone H1<sup>o</sup> Human, Recombinant with 1 µg of φX174 RF I (suprecoiled) plasmid DNA for 4 hours at 37°C resulted in < 5.0% conversion to RF II form (nicked circle) as determined by agarose gel electrophoresis.

**Protein Sequence:** TENSTSAPAAKPKRAKASKK STDHPKYSDMIVAAIQAEKNRAGSSRQSIQKYIKSH YKVGENADSIKLSIKRLVTTGVLKQTKGVGASGS FRLAKSDEPKKSVAFKKTKEIKKVATPKKASKPKK AASKAPTKKPKATPVKKAKKLAATPKKAKPKTV KAKPVKASKPKKAKPVKPKAKSSAKRAGKKK (Genbank accession number: P07305)

**Endonuclease Assay:** Incubation of a 50 µl reaction containing 10 µg of Histone H1<sup>o</sup> Human, Recombinant with 1 µg of φX174 RF I (suprecoiled) plasmid DNA for 4 hours at 37°C resulted in < 5.0% conversion to RF II form (nicked circle) as determined by agarose gel electrophoresis.

**Protein Sequence:** TENSTSAPAAKPKRAKASKK STDHPKYSDMIVAAIQAEKNRAGSSRQSIQKYIKSH YKVGENADSIKLSIKRLVTTGVLKQTKGVGASGS FRLAKSDEPKKSVAFKKTKEIKKVATPKKASKPKK AASKAPTKKPKATPVKKAKKLAATPKKAKPKTV KAKPVKASKPKKAKPVKPKAKSSAKRAGKKK (Genbank accession number: P07305)

**References:**

1. van Holde, K.E. (1989) *Chromatin*, 1–497
2. Marzluff, W.F., et al. (2002) *Genomics*, 80, 487–497
3. Pehrson, J.R. and Cole, R.D. (1982) *Biochem.*, 21, 456–460
4. Gill, S.C. and von Hippel, P.H. (1989) *Anal. Biochem.*, 182, 319–326
5. Pace, C.N. et al. (1995) *Protein Science*, 4, 2411–2423
6. Qing, X. et al. (2010) *Biochemistry*, 49, 5588–5599.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

**References:**

1. van Holde, K.E. (1989) *Chromatin*, 1–497
2. Marzluff, W.F., et al. (2002) *Genomics*, 80, 487–497
3. Pehrson, J.R. and Cole, R.D. (1982) *Biochem.*, 21, 456–460
4. Gill, S.C. and von Hippel, P.H. (1989) *Anal. Biochem.*, 182, 319–326
5. Pace, C.N. et al. (1995) *Protein Science*, 4, 2411–2423
6. Qing, X. et al. (2010) *Biochemistry*, 49, 5588–5599.



NEW ENGLAND BIOLABS® is a registered trademark of New England Biolabs, Inc.

This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.