

# 100 bp DNA Ladder



1-800-632-7799  
info@neb.com  
www.neb.com



N3231S 093120514051

## N3231S

100 gel lanes (50 µg) Lot: 0931205 Exp: 5/14  
500 µg/ml Store at -20°C

1.5 ml Gel Loading  
Dye, Blue (6X) Store at 25°C

**Description:** A number of proprietary plasmids are digested to completion with appropriate restriction enzymes to yield 12 bands suitable for use as molecular weight standards for agarose gel electrophoresis. The digested DNA includes fragments ranging from 100–1,517 base pairs. The 500 and 1,000 base pair bands have increased intensity to serve as reference points.

Supplied in: 10 mM Tris-HCl (pH 8.0), 1 mM EDTA

**Reagents supplied:**  
6X Gel Loading Dye, Blue

**1X Gel Loading Dye, Blue:**  
2.5% Ficoll-400  
11 mM EDTA  
3.3 mM Tris-HCl (pH 8.0@25°C)  
0.017% SDS  
0.015% bromophenol blue

**Preparation:** The double-stranded DNA is digested to completion with appropriate restriction enzymes, phenol extracted and equilibrated to 10 mM Tris-HCl (pH 8.0) and 1 mM EDTA.

**Usage Recommendation:** We recommend loading 0.5 µg of 100 bp DNA Ladder diluted in sample buffer. The 100 bp DNA Ladder was not designed for precise quantification of DNA mass, but can be used for approximating the mass of DNA in comparably intense samples of similar size. The approximate mass of DNA in each of the bands in our 100 bp DNA Ladder is as follows (assuming a 0.5 µg loading):


Fragment	Base Pairs	DNA Mass
1	1,517	45 ng
2	1,200	35 ng
3	<b>1,000</b>	<b>95 ng</b>
4	900	27 ng
5	800	24 ng
6	700	21 ng
7	600	18 ng
8	<b>500,517</b>	<b>97 ng</b>
9	400	38 ng
10	300	29 ng
11	200	25 ng
12	100	48 ng

**Notes:** All fragments have 4-base, 5' overhangs that can be end labeled using T4 Polynucleotide Kinase (NEB #M0201) or filled-in using DNA Polymerase I, Klenow Fragment (NEB #M0210) (1). Use  $\alpha$ -[<sup>32</sup>P] dATP or  $\alpha$ -[<sup>32</sup>P] dTTP for the fill-in reaction.

100 bp DNA Ladder is stable for at least 3 months at 4°C.

For long term storage store at -20°C. If samples need to be diluted, use TE or other buffer of minimal ionic strength. DNA may denature if diluted in H<sub>2</sub>O.

100 bp  
DNA Ladder  
visualized by  
ethidium bromide  
staining on a 1.3%  
TAE agarose gel.  
Mass values are for  
0.5 µg/lane



(see other side)

CERTIFICATE OF ANALYSIS

# 100 bp DNA Ladder



1-800-632-7799  
info@neb.com  
www.neb.com



N3231S 093120514051

## N3231S

100 gel lanes (50 µg) Lot: 0931205 Exp: 5/14  
500 µg/ml Store at -20°C

1.5 ml Gel Loading  
Dye, Blue (6X) Store at 25°C

**Description:** A number of proprietary plasmids are digested to completion with appropriate restriction enzymes to yield 12 bands suitable for use as molecular weight standards for agarose gel electrophoresis. The digested DNA includes fragments ranging from 100–1,517 base pairs. The 500 and 1,000 base pair bands have increased intensity to serve as reference points.

Supplied in: 10 mM Tris-HCl (pH 8.0), 1 mM EDTA

**Reagents supplied:**  
6X Gel Loading Dye, Blue

**1X Gel Loading Dye, Blue:**  
2.5% Ficoll-400  
11 mM EDTA  
3.3 mM Tris-HCl (pH 8.0@25°C)  
0.017% SDS  
0.015% bromophenol blue

**Preparation:** The double-stranded DNA is digested to completion with appropriate restriction enzymes, phenol extracted and equilibrated to 10 mM Tris-HCl (pH 8.0) and 1 mM EDTA.

**Usage Recommendation:** We recommend loading 0.5 µg of 100 bp DNA Ladder diluted in sample buffer. The 100 bp DNA Ladder was not designed for precise quantification of DNA mass, but can be used for approximating the mass of DNA in comparably intense samples of similar size. The approximate mass of DNA in each of the bands in our 100 bp DNA Ladder is as follows (assuming a 0.5 µg loading):


Fragment	Base Pairs	DNA Mass
1	1,517	45 ng
2	1,200	35 ng
3	<b>1,000</b>	<b>95 ng</b>
4	900	27 ng
5	800	24 ng
6	700	21 ng
7	600	18 ng
8	<b>500,517</b>	<b>97 ng</b>
9	400	38 ng
10	300	29 ng
11	200	25 ng
12	100	48 ng

**Notes:** All fragments have 4-base, 5' overhangs that can be end labeled using T4 Polynucleotide Kinase (NEB #M0201) or filled-in using DNA Polymerase I, Klenow Fragment (NEB #M0210) (1). Use  $\alpha$ -[<sup>32</sup>P] dATP or  $\alpha$ -[<sup>32</sup>P] dTTP for the fill-in reaction.

100 bp DNA Ladder is stable for at least 3 months at 4°C.

For long term storage store at -20°C. If samples need to be diluted, use TE or other buffer of minimal ionic strength. DNA may denature if diluted in H<sub>2</sub>O.

100 bp  
DNA Ladder  
visualized by  
ethidium bromide  
staining on a 1.3%  
TAE agarose gel.  
Mass values are for  
0.5 µg/lane



(see other side)

CERTIFICATE OF ANALYSIS

Due to the limitations of the acrylamide gel technology, one or two extra bands may be visible on the DNA ladders when run on a polyacrylamide gel.

**Suggested protocol for loading a sample:**

The following protocol is recommended for a 5 mm wide lane.

1. Prepare loading mixture:

Distilled water	4 $\mu$ l
6X Blue Loading Dye	1 $\mu$ l
DNA Ladder	1 $\mu$ l
Total volume	6 $\mu$ l

2. Mix gently
3. Load onto the agarose gel

**Note:** The components of the mixture should be scaled up or down, depending on the width of the agarose gel.

Due to the limitations of the acrylamide gel technology, one or two extra bands may be visible on the DNA ladders when run on a polyacrylamide gel.

**Suggested protocol for loading a sample:**

The following protocol is recommended for a 5 mm wide lane.

1. Prepare loading mixture:

Distilled water	4 $\mu$ l
6X Blue Loading Dye	1 $\mu$ l
DNA Ladder	1 $\mu$ l
Total volume	6 $\mu$ l

2. Mix gently
3. Load onto the agarose gel

**Note:** The components of the mixture should be scaled up or down, depending on the width of the agarose gel.

**Reference:**

1. Sambrook, J., Fritsch, E. F. and Maniatis, T. (1989). *Molecular Cloning: A Laboratory Manual*, (2nd ed.), (pp. 10.51–10.67). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.

**Reference:**

1. Sambrook, J., Fritsch, E. F. and Maniatis, T. (1989). *Molecular Cloning: A Laboratory Manual*, (2nd ed.), (pp. 10.51–10.67). Cold Spring Harbor: Cold Spring Harbor Laboratory Press.